This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

WORLD INTELLECTUAL PROPERTY ORGANIZATI'. International Bureau

US



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5: C12N 15/12, 15/13, 15/62 C12P 21/08, G01N 33/48, 33/534 A61K 37/02, 39/00

(11) International Publication Number: A1

WO 93/23537

(43) International Publication Date:

25 November 1993 (25.11.93)

(21) International Application Number:

PCT/US93/04338

(22) International Filing Date:

7 May 1993 (07.05.93)

(30) Priority data:

07/881,109

8 May 1992 (08.05.92)

(72) Inventors; and

(75) Inventors/Applicants (for US only): HUSTON, James, S. [US/US]; 5 Drew Road, Chestnut Hill, MA 02167 (US). KECK, Peter, C. [US/US]; 50 Dolan Road, Millbury, MA 01527 (US).

(74) Agents: BROOK, David, E. et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173

(60) Parent Application or Grant (63) Related by Continuation

Filed on

07/881,109 (CIP) 8 May 1992 (08.05.92)

(71) Applicant (for all designated States except US): CREATIVE BIOMOLECULES [US/US]; 35 South Street, Hopkinton, MA 01748 (US).


(81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: CHIMERIC MULTIVALENT PROTEIN ANALOGUES AND METHODS OF USE THEREOF





(57) Abstract

A chimeric immunoglobulin (Ig) Superfamily protein analogue comprising one or more polypeptide chains forming a βbarrel domain having more than one biologically active binding site is disclosed. Also disclosed are methods of designing and producing chimeric multivalent Ig Superfamily protein analogues and compositions comprised of these analogues and methods of use thereof.

> Atty Dock. No: Serial No.: Reference:

6750-018-999 09/831,631

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

		co	Company	MR	Mauritania
АŦ	Austria	FR	France		
ΑU	Australia	GA	Gabon	MW	Malawi
88	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinca	NO	Norway
BF.	Burkina Faso	GR	Greece	NZ	New Zealand
BC	Bulgaria	HU	Hungary	PL,	Poland
BJ		ie ie	Ireland	PT	Portugal
BR	Brazil	1T	ltaly	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SK	Slovak Republic
CI	Côte d'Ivoire	KZ	Kazakhstan	SN	Senegal
CM	Canterout	1.1	Liechtenstein	SU	Soviet Union
CS.	Czechoslovakia -	LK	Sri Lunka	TD	Chad
cz	Czech Republic	1.0	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	UA	Ukraine
DK	Denmark	MG	Madagascar	US	United States of America
		· ML	Mali	VN	Vict Nam
	•				
ES FI	Spain Finland	.MN	Man Mongolia	***	A ICC LATIN

25

1

CHIMERIC MULTIVALENT PROTEIN ANALOGUE METHODS OF USE THEREOF

Description

Background of the Invention

The Immunoglobulin (Ig) Gene Superfami. comprised of numerous cell surface and solul molecules that mediate recognition, adhesion functions in vertebrates. (Abbas, A.K. et a AND MOLECULAR IMMUNOLOGY, p.144 (1991)). Me 10 the Iq Superfamily have an evolutionary rela and share significant amino acid sequence as structural similarities. (Williams, A. F. a) A. N., IMMUNOGLOBULIN GENES, p.372 (1989)). criteria for membership within the family a: 15 sequence homology with Ig or Ig-related pol; domains, which are approximately 70-110 ami) residues long, and 2) key structural feature include the polypeptide domains comprised o arrangement of two β -sheets, each made up of 20 five anti-parallel β -strands of five to ten residues. (Abbas, A.K., et al., CELLULAR AN IMMUNOLOGY, pp. 144-145 (1991)).

The Ig Superfamily domains are classif either variable (V) or constant (C) based o characteristics of the β -strands within the sandwich. (Abbas, A. K., et al., CELLULAR A IMMUNOLOGY, p. 144 (1991)). For example, i of Superfamily molecules, the immunoglobuli

15

20

25

30

domains are at the amino-terminal ends of separate "heavy" (H) and "light" (L) chains, succeeded in the polypeptide chains by constant (C) domains. Thus, in an immunoglobulin, a V domain is defined as either V_H or V_L (Figure 1). In the T cell receptor, V and C refer to Ig-like variable and constant domains which are comprised of polypeptide α and β chains. In other superfamily molecules, V and C domains may be comprised of γ , δ , or ϵ chains.

In the Ig Superfamily, the polypeptide chains that comprise the V regions, associate to form ligand binding sites. For example, in the immunoglobulin molecule, the $\boldsymbol{V}_{\boldsymbol{H}}$ and $\boldsymbol{V}_{\boldsymbol{L}}$ domains associate to form the variable fragment (Fv) region which comprises the antibody binding site. The Fv region includes both scaffold-like regions, termed framework regions (FRs), and regions of hyper-variability, termed complementarity-determining regions (CDRs). It is the CDRs that contribute to the unique antigen specificity of immunoglobulins. (Abbas, A. K., et al., CELLULAR AND MOLECULAR IMMUNOLOGY, p.45 (1991)). Under special circumstances, the Fv region has been proteolytically dissected from its parent Ig to yield a variable-region fragment (Fv fragment) that is comprised of two noncovalently associated domains $(V_H \cdot V_L)$, a heterodimer). This heterodimeric Fv fragment can further dissociate into single V_L and V_H domains. (Huston, J.S. et al. Meth. Enzymol. 203:46-88 (1991)).

Recently, protein engineering methods have been used to link V_H and V_L chains, creating a functional single-chain Fv (sFv), which has a solitary antibody binding site that does not dissociate into single domains at low concentrations. (Huston, J.S., et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (August 1988)).

-3-

In this approach, the genes encoding V_H and V_L domains of a given antibody are connected at the DNA level by an appropriate nucleotide sequence, and on translation, this gene forms a single polypeptide chain with a peptide linker bridging the two variable domains. (Huston, J.S., et al., Meth. Enzymol., 203:46-88 (1991)).

Summary of the Invention

5

25

30

The present invention relates to a chimeric immunoglobulin (Ig) Superfamily protein analogue having 10 more than one biologically active binding site. Hereinafter, the term multivalent will be used to describe these multiple binding sites. The chimeric multivalent Ig Superfamily protein analogue, hereinafter referred to as a CHI-protein, or χ -protein, 15 is comprised of one or more polypeptide chains forming a β -barrel domain. A single β -barrel domain may comprise a chimeric protein binding domain with more than one binding site. Alternatively, more than one β barrel domain, such as the V_L and V_H β -barrel domains 20 in immunoglobulins, may combine to form a larger concentric β -barrel domain having more than one binding site.

The β -barrel domain(s) comprising the binding regions has amino acid sequence and structural homology with variable regions of molecules related to the Ig Superfamily of molecules. Specifically, the binding sites on the χ -protein are comprised of hypervariable regions derived from molecules related to the Ig Superfamily of molecules. The Ig Superfamily includes immunoglobulins, cell surface antigens, such as T lymphocyte antigens, and cell surface receptors, such as immunoglobulin Fc receptors.

20

25

30

35

In a preferred embodiment, the hyperva regions are complementarity-determining reg derived from the antigen binding sites of immunoglobulins. In this embodiment, the m protein analogue comprises one or more poly chains forming a β -barrel domain containing interspersed between framework regions (FRs CDRs define one antigen binding site. This protein analogue also has one or more addit antigen binding sites spliced into the FRs barrel domain.

In one embodiment, the χ -protein will single polypeptide chain forming a β -barrel In another embodiment the χ -protein will cosingle polypeptide chain comprised of two polypeptide chains, connected by a polypeptide linker so distance between the C-terminus of one chain terminus of the other chain forming a β -barrel in yet another embodiment, the χ -protein with two polypeptide chains with two non-covaler associated chains forming a β -barrel domain of the above embodiments, the polypeptide contains the polypeptide contains a β -barrel domain with two or more sites.

The invention also relates to the amir sequences that encode the χ -protein, the DN that encode the amino acid residues forming protein, and expression vectors comprising of expressing the DNA sequences. The invent relates to methods of producing the χ -prote

The invention further relates to compositions are comprised of χ -proteins and methods of These compositions are comprised of χ -proteins about two biologically active binding sites and in a variety of therapeutic and diagnostic

These compositions include, but are not limited to, χ protein biosensors which undergo a conformational
change when a ligand is bound to one binding site such
that the affinity of the second binding site is
modified; and χ -proteins having one binding site
reactive with a tissue-specific ligand and the second
binding site reactive with radioactive ions, radioopaque substances, cytotoxic substances, cytotoxic
effector cells (e.g. cytotoxic T cells) drugs, or
catalytic substances. These compositions also include
a "biochip" comprised of a two-dimensional array of
aggregated χ -protein biosensors, such as in a LangmuirBlodgett film, to make a functionalized membrane useful
for layers of molecular gates for computers and the
like.

5

10

15

20

25

30

35

The utility of binding proteins having two independent binding sites of different specificity for the treatment or control of tumors, virus infected cells, bacteria and other pathogenic states has been recognized (Segal, D.M. and Snider, D.P., Chem. Immunol. 47:179-213 (1989)). Bispecific binding proteins have been produced by crosslinking two or more dissimilar but intact antibodies with a chemical agent (heteroantibodies); crosslinking antibody fragments; linking two single chain antibodies; or fusing a single chain antibody to an effector molecule (Segal et al. U.S. 4,676,980), Tai, M., et al. Biochem. 29:8024-8030 (1990).

However, despite considerable application to medical research, these previous attempts to produce bispecific binding proteins suffer from the difficulty of complex purifications and large molecular size. For example, each binding region of an IgG is at least 50 kD, so that a conventional crosslinked, bispecific heteroantibody can range in mass from 100-150 kD to as

20

25

30

much as 300 kD, if two intact IgG antibodies are cross-linked. Even the single chain antibody has a mass of approximately 26 kD so that a single polypeptide chain comprising two separate binding regions, each comprised of a single chain antibody, has a mass of approximately 50 kD.

The recombinant-engineered x-proteins of the present invention have significant advantages over conventionally crosslinked antibodies, or even the single-chain Fv. The proteins of the present invention can be custom-designed to bind specific ligands and cell surface receptors with affinity or specificity. These custom-designed multivalent binding proteins can be smaller and more compact in size than intact hetero-bispecific antibodies, Fab' antibody binding fragments or bispecific sFv-sFv constructs.

These χ -proteins can also be less immunogenic thereby reducing the likelihood of immune reactions to such therapeutic compositions. For example, in the intact immunoglobulin molecule, the bottom loops of the variable region are sterically protected from recognition resulting in an immune response. However, when disassociated from the protecting constant region, as in the Fv, the exposed bottom loop region potentially becomes antigenic. In a χ -protein, this bottom loop region is no longer exposed and thus significantly reduces the likelihood of an immune response. Furthermore, due to their smaller size, the x-proteins can have enhanced stability when administered intravenously as they will be less susceptible to proteolysis by endogenous proteases than larger, multi-domain proteins.

10

15

20

25

30

Brief Description of the Drawings

The foregoing features and advantages invention will be apparent from the follow: particular description in the following dratext.

Figure 1 is a schematic representation typical Ig Superfamily molecule, an immunoc depicting the variable (V) and constant (C)

Figure 2A is a schematic representation depicting the relative positions of the top and bottom loops (BL) of the folded heavy a chains. The TLs typically form the CDRs of the BLs are typically adjacent to the C receivithin an intact immunoglobulin. The BLs of loops suitable for splicing on the second be

Figure 2B shows an alignment of V_L and acid sequences for which three dimensional are known (SEQ ID NOS: 1-9 for V_L and SEQ I for V_H). The alignment of these sequences the structural homology that exists between especially in the β -strand regions. Region sequence corresponding to structural region β -strand (IS), outer- β -strand (OS), top low bottom loops (BL) are identified. The number corresponds to structural position, especially and IS regions.

Figure 3A is a stereo figure depicting of the McPC 603 Fv structure wherein the to one (right side up) structure are superimple bottom loops of the other (upside down) structure, only the top loops (ribbons) and (line trace) are shown.

10

20

25

30

Figure 3B is a stereo figure depicting the alignment of H2 with the N- and C-terminal strands of the inverted Fv structure.

Figures 4A-4G depict in stereo the positions of the native CDRs and the CDRs spliced into the McPC 603 bottom loops to form a additional binding site (χ -site). Corresponding native and χ -site CDR loops are highlighted as ribbons.

Figure 5 depicts the stereo comparison of the χ -protein comprised of native McPC603 with McPC603 CDRs spliced into the BLs of the native McPC603.

Figure 6 depicts the splice points used when H3 of a second McPC603 is spliced into H BL2 of the native McPC603 sFv. The ribbon follows the spliced trace.

Figure 7 depicts the splice points used when the L3 loop is spliced into the L LB2 of the native McPC603.

Figure 8 depicts the splice points used when the H1 loop is spliced into H BL4 of the native McPC603.

Figure 9 depicts the splice points used when the L1 loop is spliced into L BL4 of the native McPC603.

Figure 10 depicts the splicing of the H2 loop into the C-terminus of $\ensuremath{V_{H}}$ of native McPC603.

Figure 11 depicts the splicing of the L2' loop onto the C-terminus of V_L of native McPC603.

Figure 12 shows the final amino acid sequence of the χ -protein comprised of two McPC603 binding sites as constructed according to Examples 1 and 2 (SEQ ID NO: 17).

Figures 13 A and B show the alignment of consensus sequences for the various sequence classes in the Kabat et al. compendium with the alignment of sequences of known structure for the V_H (Figure 13A) and V_L (Figure 13B) chains; (Kabat, E.A., SEQUENCES OF PROTEINS OF

15

20

25

30

IMMUNOLOGICAL INTEREST Vols. I-III, U.S. Dept. Health and Human Services, NIH Pub. No. 91-3242 (1991)). The residue preference list in Table 2 is derived from this figure. The first group of sequences are as in Figure 2B (SEQ ID NOS: 1-16). The second group of sequences are consensus sequences from the groupings in Kabat et al. (Figure 13A, SEQ ID NOS: 18-28) and Figure 13B, SEQ ID NOS: 31-38). The third group of sequences are known sequences for mouse immunoglobulins 2610 and GLOOP4. (Figure 13A, SEQ ID NOS: 29-30 and Figure 13B, SEQ ID

10 (Figure 13A, SEQ ID NOS: 29-30 and Figure 13B, SEQ ID NOS: 39-40). The line labeled KABAT indicates the boundaries of framework and CDR regions as defined in Kabat et al. ("-" indicate alignment gaps).

Figure 14 shows the sequences of χ -protein constructs as constructed according to Examples 2 and 3 (SEO ID NOS: 41-44).

Figure 15A shows the nucleic acid and amino acid sequences of the χ -1 protein, with double dashes indicating the D1.3H1 χ -loop insertion in the 26-10 sFv (SEQ ID NOS: 45 AND 46, respectively).

Figure 15B shows the nucleic acid and amino acid sequences of the χ -2 protein, with double dashes indicating the D1.3H1 and H2 χ -loop insertions in the 26-10 sFv (SEQ ID NOS: 46 and 47, respectively).

Figure 16A shows the SDS polyacrylamide gel electrophoresis of the χ -1 protein purified by ouabain-Sepharose affinity chromatograph.

Figure 16B shows the SDS polyacrylamide gel electrophoresis of the $\chi-2$ protein purified by ouabain-Sepharose affinity chromatography.

Figure 17 shows the elution profiles of the $\chi - 2$ protein and the 26-10 sFv from a Superdex 75 column.

Detailed Description of the Invention

20

25

30

The present invention relates to a chim multivalent immunoglobulin (Ig) Superfamily analogue, hereinafter referred to as a χ-pro comprising one or more polypeptide chains for barrel domain. The β-barrel domain, contain hypervariable regions (hereinafter called complementarity-determining region-like (CDI region)) and structural regions (hereinafter framework-region-like (FR-like region)). The regions define ligand binding sites. Additing the protein has at least one more ligand binding segment spliced into the FR-like regions of barrel domain.

The Ig Superfamily molecules show sign: amino acid sequence homology within their $oldsymbol{eta}$ (Williams, A.F. and Barclay, A.N. domains. IMMUNOGLOBULIN GENES p. 362 (1989)). The a sequences of the amino terminal domains are variable (V) regions and the more conserved of the remainder of the chain, termed the c (Figure 1) (Abbas, A. K., et al., region. MOLECULAR IMMUNOLOGY p. 45 (1991)). The am sequences of both the V and C regions of Su molecules are formed on two different polyp chains, the heavy (H) chain and the light (immunoglobulins, or the α , β , γ , δ , or ϵ ch other Ig Superfamily molecules. These chai into V and C regions. For example, each H immunoglobulin molecule folds into a $V_{\scriptscriptstyle H}$ dom adjacent C_H domain and each L chain folds i domain with an adjacent C_L . Each chain has successive constant regions (Figure 1).

The V region contains highly variable stretches of sequence called the hypervaria

More conserved stretches of sequence are called structural regions. In immunoglobulins, the hypervariable regions are called complementaritydetermining regions (CDRs) and the structural regions are called framework regions (FRs). Three CDRs of each V_H, and three CDRs of each V_L combine in a unique three-dimensional structure to form the antigen or These CDRs determine the ligand ligand binding site. specificity of the protein. (Abbas, A. K., et al., CELLULAR AND MOLECULAR IMMUNOLOGY p.143 (1991)). Hereinafter, the hypervariable regions of all Ig Superfamily molecules will be called CDR-like and all conserved regions will be called FR-like (or CDRs and FRs in the specific case of an immunoglobulin molecule).

5

10

15

20

25

30

The Ig Superfamily members also show significant homology in the structural three-dimensional features of their V and C domains. This structural feature is known as the Ig-fold. (Williams, A.F. and Barclay, A.N., IMMUNOGLOBULIN GENES p. 362 (1989)). The Ig-fold consists of a sandwich of two β -sheets constructed from anti-parallel β -strands, each strand containing five to ten amino acid residues. The V domain differs from the C domain by an extra pair of β -strands in the middle of the V domain. (Williams, A.F. and Barclay, A.N., IMMUNOGLOBULIN GENES p. 362 (1989)). For example, the V domains or C domains of an immunoglobulin, associate in pairs, such as $V_H \cdot V_L$. Each β -barrel domain, when associated in such a pair, forms two concentric β barrels, with the CDR loops connecting anti-parallel β strands of the inner barrel.

Members of the Ig Superfamily include, but are not limited to, Immunoglobulins, T cell Receptor Complex, Major Histocompatibility Complex Antigens, β_2

10

15

20

25

30

Microglobulin-associated Antigens, T Lymphocyte
Antigens, Growth Factor Receptors, and Neural Cell
Adhesion Molecules (NCAM). (Williams, A.F. and Barclay,
A.N., IMMUNOGLOBULIN GENES p. 362 (1989)).

Each ligand binding site of the χ -protein is comprised of the CDR-like region derived from molecules of the Ig superfamily. For example, a ligand binding site could be comprised of the CDRs derived from an immunoglobulin molecule whose ligand is an antigen. Alternatively, the ligand binding site could be comprised of CDR-like regions from a receptor molecule such as the T cell receptor whose ligand is the antigen-major histocompatibility complex (MHC) molecule which, upon binding to its receptor, initiates T cell activation.

In particular, the present invention relates to the immunoglobulins of the Ig Superfamily. In natural immunoglobulins, the antibody combining site is formed by CDRs of the V_H and V_L variable domains within the Fv (variable region consisting of noncovalently associated V_H and V_L or $V_{H^{\circ}}V_L$), as shown in Figure 1. In addition to the CDRs determining binding specificity, maintenance of the tertiary structure is also necessary for biological activity. The CDRs are correctly positioned by the conserved framework regions (FRs) within the V regions, and the V region is further stabilized by the C regions of the protein. However, the minimal naturally-occurring antibody binding site is the two chain, non-covalently associated Fv.

Recently, through recombinant protein engineering, a single-chain Fv (sFv) has been constructed. In this approach, the genes encoding $V_{\rm H}$ and $V_{\rm L}$ domains of a given antibody are connected at the DNA level by an appropriate oligonucleotide linker and, on translation,

10

15

20

25

30

this gene forms a single polypeptide chain v peptide linker bridging the two variable don (Huston, J.S. et al., Meth. Enzymol. 203:46-

Isolated single domain antibodies have recently been constructed, comprised of only instead of the usual complement of six. (Water al., Nature 341: 544-546 (1989)). In sorthese single domain antibodies (i.e., VH or binding activities comparable to their parer antibodies.

In a preferred embodiment of the present invention, one of the χ -protein binding site comprised of a set of CDRs from a mouse myet such as 26-10 or McPC603 (Huston, J.S., et a Enzymol. 203: 46-88 (1991). An additional segment is spliced into the β -barrel domain spliced segment is comprised of CDRs from the a second mouse myeloma protein, which are split the bottom FR loops of the β -barrel domain, to the C-terminal ends of each V domain.

The structural basis for this invention explained with reference to Figure 2B (SEQ 16), where the typical variable region compan immunoglobulin is noted. There are threwithin each V_H and V_L which together constitute complete antibody binding site. These CDRs interspersed between FRs such that the light region is denoted FR1-L1-FR2-L2-FR3-L3-FR4 heavy chain is denoted FR1-H1-FR2-H2-FR3-H3 of these V region chains folds into a native conformation that comprises a double layer V domains may be monomeric, $(V_H \text{ or } V_L)$, or into the homodimers, $(V_H \cdot V_H \text{ or } V_L \cdot V_L)$, or into the heterodimeric FV, $(V_H \cdot V_L)$. Each of these dimerical explanations are such that the second explanation is denoted from the second explanation that comprises a double layer V domains may be monomeric, $(V_H \cdot V_L)$, or into the heterodimeric FV, $(V_H \cdot V_L)$. Each of these dimerical explanation is denoted from the second explanation that the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanation in the second explanation is denoted from the second explanat

constructed as single-chain analogues. In the single-chain Fv, these two sequences are connected in tandem with a bridging linker to make, for example, V_H -linker- V_L (V_H - V_L) or V_L -linker- V_H (V_L - V_H).

5

10

15

20

25

30

In these folded configurations of the V region polypeptide, alternating directions of the β -strands loop out at either end as they form anti-parallel strands. In each region the immunoglobulin fold has loops on top (the binding site is "on top") and on the bottom (partly in close proximity to constant domains for intact H and L chains).

On top of each V domain, four loops are present of which three are CDRs that contribute to the antigen binding site, and on the bottom, four loops are present that allow the β -strands to switch directions and fold back into the globular domain. These bottom loops are here termed BL1, BL2, BL3, and BL4, and apply to the L and H variable regions, respectively called L BL1-4 and H BL1-4 (Figure 4A). These bottom loops provide insertion sites for fusion proteins or peptide segments, which are further augmented by splicing of peptide sequence at the C-terminus of each V region. The use of these bottom loops as splice sites permits incorporation of alternate binding, catalytic or effector sites which complement the naturally present antigen binding site.

However, novel demands are made on variable region architecture in order to construct additional binding sites on the bottom of a V domain. In particular, there is a requirement that the relative orientation of the CDR-like loops (CDR or top loop symmetry) for an Fv region be reproduced to a reasonable approximation in the relative orientation of the bottom loops of the Fv (bottom loop symmetry). Whether or not this symmetry

-15-

relationship exists is not an obvious question to ask and the possibility of top-bottom loop symmetry has not been previously recognized. Nonetheless, molecular modeling and computational analysis demonstrate that such an approximation to the CDR-like loop symmetry does exist for the bottom loops of the Fv region.

5

10

15

20

25

30

The superposition of top and bottom loops involves two interrelated observations: (1) the two fixed endpoints of each top loop (CDR) must find a corresponding match with bottom loop endpoints; and (2) the polypeptide chain directionality must be maintained after bottom loop splicing (i.e., N to C directionality of the spliced CDR must be the same as the bottom loop that it replaces). The overall architecture of the Fv region appears asymmetric because the ends of the light and heavy chain β -barrels are closer together at the top than they are at the bottom. However, much of this apparent difference derives from there being one quarter of the Fv residues in CDRs on top of the framework, which fill in space that is open on the bottom.

In fact, there is sufficient correlation between top and bottom loop symmetry of the intact Fv region to make the multiple binding site molecule possible. This is discernible if one superimposes the top of one Fv (#1) with the bottom of another Fv (#2); if #1 and #2 are the same, then the correlation helps decide how to construct a multivalent Fv, whereas if #1 and #2 are different, the additional binding site (χ -site) is distinct from the native Fv binding site, and overall one thereby designs a chimeric multivalent Ig Superfamily protein analogue which has one or more sites of different specificity.

We will represent the composition of the χ -site with parentheses so that A(B) represents a χ -protein

20

30

comprising the CDR-like regions of immunogle Superfamily molecule B built into the χ -site immunoglobulin Superfamily molecule or analogical Likewise ($V_H(V_L)$) represents a V_H domain with whose loops are derived from the CDR loops domain. Consequently the designation A(B) denotes a χ -protein based on the FR and CDR having a χ -site based on the CDR loops of F the χ -site loops on the heavy chain of A around the CDR loops of the light chain B and and the two χ domains are linked by a polyplinker such that the heavy chain FR domain light chain FR domain.

As shown in Figure 3A, by applying the procedures to McPC603 for both Fv #1 and #2 following pairs of loops align:

- The TL4 loops (L3 and H3) of Fv#1 can very closely with the BL2 loops of Fv# the alignment of these two sets of loo additional alignments become apparent.
- 2. The TL1 loops (L1 and H1) are approxim superimposed on the respective BL4 loo
- In addition, as shown in Figure 3B, the TL2 loops (L2 and H2) are approximal superimposed on the N-terminal and C-t strands of the respective β-barrel domestic formula in the strands of the respective β-barrel domestic formula in Figure 3B, the strands of the respective β-barrel domestic formula in Figure 3B, the strands of the N-terminal and C-t strands of the respective β-barrel domestic formula in Figure 3B, the strands in Figure 3B, the s

In this structural alignment, all of t fundamental design criteria for splicing a binding site onto the bottom of V, Fv, or s are satisfied: the proper chain direction

PCT/US93/04338 WO 93/23537

superpositions are found for the TL4 - BL2 pair, the TL1 - BL4 pair, and the ends of the TL2 loops relative to the N- and C-terminal strands of the β -barrel region.

5

10

20

25

30

This construction is consistent to a good approximation with the natural geometry for all CDR loops, thereby generating a ligand binding site on the bottom of the β -barrel domain similar to the "source" Fv binding site. Similarly, an additional ligand binding site may be built on the separate noncovalently linked V domains of a heterodimeric Fv (i.e., $V_{H} \cdot V_{L}$). Alternatively, as it has recently been shown that the full set of 6 CDRs is not necessary for ligand binding, partial binding sites (i.e., fewer 15 'than 6 CDRs) may be assembled on a single V domain. Thus, a χ -protein could comprise a single β -barrel domain with two ligand binding sites, each binding site with as few as one CDR and still exhibit binding activity.

H BL1 or L BL1 loop replacement could also be useful, as they are in proximity to the rest of the binding site. However, their peptide chain directionality is opposite to what would be needed to correctly splice in H2 and L2 loops. Nonetheless, such loops could be devised de novo by design or mutagenesis to facilitate such replacement.

Furthermore, in most Fv regions, the H or L BL3 loops are located on the sides of the V domains. Although these loops are not contiguous with the rest of the additional binding site, ancillary loops could be attached at such sites, thereby providing the addition of a "ligand-like" surface feature for recognition by the appropriate receptor, thus, forming a χ -protein with more than two binding sites. A single

10

15

20

25

30

35

substitution of only one loop, with an appropriate peptide, could provide a means to anchor the binding protein to a particular receptor. Moreover, the H or L BL3 (or H or L TL3) loops could provide the means to crosslink single domain χ -proteins into aggregate sheets to form two dimensional arrays of χ -proteins.

CDR-like region sequences of any of the proteins from the Ig Superfamily showing the requisite sequence and structural homology can be spliced into the bottom loops of the source Fv. It should be noted that CDR-like region sequences from other Ig Superfamily molecules can also replace all or a part of the native CDR-like region of the χ -protein. Thus a χ -protein can comprise the CDR-like regions from molecule A, the FR-like regions from molecule B and the χ -site having CDR-like regions from molecule C spliced in.

As discussed in detail in Examples 1 and 2, using the Fv derived from mouse myeloma IgA antibody, McPC 603, a second McPC 603 binding site can be constructed on the bottom of the source Fv. Although the Fv region of McPC 603 is exemplified, as shown in Example 3, it is reasonable to splice in a CDR-like region from other Ig Superfamily proteins, due to the sequence and structural homology that exists among these proteins. (Figure 2B (SEQ ID NOS: 1-16)).

When CDR-like sequences are spliced into the lower FR-like loops of the source Fv, it is important to preserve those FR-like residues that are critical for maintaining the proper folding. These critical FR-like residues are located in the stems of the loops and underlying the CDR-like regions. Three criteria govern the selection of the proper splice points. First, as discussed above, the need to preserve critical FR-like residues; second, the desire to incorporate as much of the CDR-like sequence as possible; and third, the

-19-

practical need to switch from one backbone to the other at points where the alpha carbons of the two chains are reasonably well aligned. The last requirement is necessary in order for the spliced loops to maintain their native, biologically active conformation.

5

10

20

25

30

As described in detail in Examples 1 and 2, the crystallographically determined coordinates of a mouse myeloma protein structure, such as McPC603, are visually displayed using a suitable computer graphics system. This display of the protein in threedimensions, can be spatially rotated, turned or twisted as necessary to view the polypeptide chains or peptide backbone of the Fv region. Using the graphics program, substitutions, additions or deletions can be made to the structure. These modifications are energy minimized (optimized) to account for steric hindrance, bond lengths, bond angles and energy constraints so as to maintain the critical tertiary structure necessary for biological binding activity. Thus, in a step by step process involving modification of the polypeptide chain and successive cycles of refinement calculations for each modification, a three-dimensional model of the χ -protein, with two distinct binding sites, is built.

As described in Examples 1 and 2, the model used was McPC603 sFv constructed as $V_H - V_L$. However, sFv proteins constructed as $V_L - V_H$, $V_H - V_H$, or $V_L - V_L$ are also intended to be encompassed by this invention. Two chain Fv regions (i.e., non-covalently associated V_H and V_L domains, $(V_H \cdot V_L)$, as well as single domain antibodies $(V_H$ or $V_L)$ are also intended to be encompassed by this invention. In addition, any other Ig Superfamily molecule having the requisite sequence and structural homology are also intended to be encompassed by this invention.

-20-

Alternatively, since the sequence would be known for the Ig superfamily CDR-like regions of interest, alignment procedures that relate tertiary to primary structure are useful to predict successful χ -proteins. Figures 13A (SEQ ID NOS: 10-16 and 18-30) and B (SEQ ID NOS: 1-9 and 31-40) depicts the amino acid residue alignment of Fv regions from mouse myeloma antibodies and other Ig superfamily molecules. As described in detail in Example 3, this type of sequence alignment allows approximate choice of splice points without having a three-dimensional crystallographic structure. The tertiary structures of framework regions are highly conserved, thus allowing reliable three-dimensional models to be predicated on sequence alignment methods.

5

10

15

20

25

30

However, to correctly splice "target" CDRs-like regions into the source Fv, some modifications to the source FR-like region sequence may be necessary.

Moreover, it may be desirable to modify the native CDR-like sequence which is spliced into the source Fv, or the source Fv itself, to modify binding affinity or specificity. Such modifications are intended to be encompassed by the subject invention.

For example, one or more amino acid residues can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs, such as the nonpolar, polar, and positively or negatively charged amino acids.

The structure may be modified by deletions, additions, substitutions and insertions of one or more amino acids which do not substantially detract from the desired functional properties of the χ -protein.

35 Naturally occurring allelic variations and

-21-

modifications are included within the scope of this invention so long as the variation does not substantially reduce the ability of the χ -protein to bind its ligand.

5

10

15

20

25

30

Based on the method described in Example 3, a χ -protein has been partially constructed incorporating two loops of an anti-lysozyme monoclonal antibody in the 26-10 sFv, as described in detail in Example 4. As shown in Figure 15A and 15B (SEQ ID NOS: 45-48), constructions have been made which incorporate the H1 and H2 loops of the D1.3 anti-lysozyme monoclonal antibody in the appropriate χ sites at the bottom of 26-10 sFv, where they are designated H1' and H2'. design process has involved successive addition of χ loops, so that there are constructions with H1' alone (χ -1, Figure 15A), and H1' + H2' together (χ -2, Figure In constructing χ -1 and χ -2, the only deviations from the corresponding partial constructs described in Example 3 (Figure 14, 2610(D1.3) sequence) are the following: (1) the isoleucine at the N-terminal end of H2' at V_H residue 112 is a valine in χ -2 due to limitations imposed by the restriction sites that were utilized, and (2) in both χ -1 and χ -2, the linker is (Ser₄Gly)₃Ser which reasonably confers better solubility properties on the proteins than that noted in Figure 14, Ala₅(Gly₄Ser)₃. The stepwise incorporation of the remaining D1.3 CDRs (H3', L1', L2', and L3') can be accomplished by the same procedure.

As each CDR loop was inserted in the χ -site, the cloned proteins were expressed in <u>E. coli</u> and refolded. The recovery of ouabain binding capacity by these χ -proteins demonstrates that the insertion of these χ -CDRs was compatible with proper refolding of the 26-10(D1.3) sFv polypeptide chain, thereby generating a

10

15

20

25

30

26-10 digoxin binding site in each case. (Ouabain is a digoxin analogue with K_a for association with 26-10 sFv of about $10^7~\rm M^1$, and ouabain-based affinity chromatography is the preferred isolation method for digoxin binding proteins.

Thus, experimental results support the rationale of inserting CDR loops in place of turns in B-sheet structure or near the C-terminal ends of V-regions. The maintenance of V region integrity after insertion of χ -loops is a fundamental premise for the construction of these dual binding site x-proteins. The ability to make a biologically-active xsFv protein (i.e., a χ -protein that retains its binding activity due to proper conformation) with 2 inserted loops (here comprising 24 CDR residues) is a remarkable result, in that the phage display manipulation of sFv binding sites may be used in entirely novel ways to genetically convert these added χ -site residues into virtually any desired antigen specificity. These results also reasonably support the anticipated ability to closely mimic a parent antibody combining site once more χ -CDRs have been incorporated.

The present invention also relates to the amino acid sequences encoding the χ -protein. Once the optimal splice points for CDR-like region insertion have been determined and shown to adequately reproduce the native site, the linear amino acid sequence can then be deduced. The χ -protein, or its modified equivalent, can then be made by recombinant DNA methods or synthesized directly by standard solid or liquid phase chemistries for peptide synthesis, or other methods well known to one skilled in the art. For example, the amino acid sequence of the χ -protein McPC603 (McPC603) shown in Figure 12 can be synthesized

by the solid phase procedure of Merrifield by semisynthesis or enzymatic or chemical combinations of appropriately modified blocks of peptides.

5

10

15

20

25

30

The present invention also relates to nucleic acid sequences, DNA and RNA, that encode the χ -protein, and expression vectors capable of expressing the χ -proteins. Preferably, the χ -proteins of the subject invention will be produced by inserting DNA encoding the desired amino acid sequence of the χ -protein into an appropriate vector/host system where it is expressed, or the gene may be used in a cell-free rabbit reticulocyte ribosomal protein synthesis system.

A variety of host/vector systems can be used to express the polypeptides of this invention, such as the one described in Example 4. Primarily, the vector system must be compatible with the host cell used. Host/vector systems include, but are not limited to, the following: bacteria transformed with bacteriophage DNA, plasmid DNA or cosmid DNA, microorganisms such as yeast containing yeast vectors; mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.) or expressing plasmids; insect cell systems infected with virus, such as baculovirus or Xenopus oocytes injected with DNA encoding the χ-protein. Other methods well known to one skilled in the art may also be used.

Any of the standard recombinant methods for the insertion of DNA into an expression vector can be used. The recombinant vector can be introduced into the appropriate host cells (bacteria, virus, yeast, mammalian cells or the like) by transformation, transduction or transfection, depending upon the host/vector system and cultured to express the polypeptides of this invention.

10

15

20

25

30

Although strategies used to produce recombinant proteins have often relied upon bacterial expression of fusion proteins followed by in vitro refolding of the protein, direct expression and secretion also may be used to produce the χ -proteins of the subject invention. Expressed fusion proteins may require purification and possible removal of the leader sequence before refolding of the χ -protein to recover binding activity. However, this may be accomplished using routine laboratory procedures. Direct expression of the χ -protein can potentially produce the protein without the leader, but still in need of refolding, whereas secretion can ideally produce native protein for subsequent isolation. (Huston, J.S., et al. Methods Enzymol., 203:46-88 (1991)).

Alternatively, fusion proteins may be the end product of expression. For example, a cytochrome b_5 tail could be fused to the χ -protein (at the gene level), to make a membrane anchor that could hold the χ -protein in a membrane or Langmuir-Blodgett film. A tail or leader sequence could also include a specific recognition sequence for enzymatic or chemical modification. Such an engineered, post-translational modification could allow specific incorporation of moieties such as biotin or phosphoinositol.

Possible refolding protocols include dilution refolding, redox refolding and disulfide-restricted refolding. (Huston, J.S., et al. Methods Enzymol. 203:46-88 (1991) the teachings of which are hereby incorporated by reference). Dilution refolding relies on the observation that fully reduced and denatured antibody fragments can refold on removal of denaturant and reducing agent with recovery of specific binding

PCT/US93/04338 WO 93/23537

activity. (Haber, E., Proc. Natl. Acad. Sci. U.S.A., 53:524 (1964).

Redox refolding utilizes a glutathione redox couple to catalyze disulfide interchange as the protein refolds into its native state. (Saxena; P. and Wetlaufer, D.B., <u>Biochem.</u>, 9:5051 (1971); (Huston, J.S., et al. Methods Enzymol. 203:46-88 (1991)).

5

10

20

25

30

Disulfide-restricted refolding involves initial formation of intrachain disulfides in the fully denatured protein. This capitalizes on the favored reversibility of antibody refolding when disulfides are kept intact. (Buckley, C. E., et al., Proc. Natl. Acad. Sci. U.S.A., 50:827 (1963); Huston, J.S., et al. Methods Enzymol, 203:46-88 (1991)). Disulfide crosslinks should restrict the initial refolding pathways available to the molecule. For chains with the correct disulfide pairing, the recovery of a native structure should be favored, while those chains with incorrect disulfide pairs must necessarily produce nonnative species on removal of denaturant.

Characterization of the multiple ligand binding sites requires that both their affinity and specificity be determined. Ideally, the measurement of binding affinity should use a thermodynamically rigorous approach such as equilibrium dialysis or ultrafiltration. In the absence of such methods. agreement between two distinct methods is desirable to reinforce the veracity of the data. Ligand binding to high affinity binding sites frequently involves very fast rates of binding and slow rates of dissociation. This characteristic makes measurement of their association constants amenable to routine immunoassay procedures, such as immunoprecipitation, radioimmunoassay and ELISA. (Huston, J.S., et al. Methods Enzymol. 203:46-88 (1991)).

35

10

20

25

30

After evaluating the binding activity of a particular model χ -protein, it may be desirable to further refine the initial χ -protein to enhance its biological activity. This refinement may be performed computationally by additional computerized modeling or it may be a "biological" refinement using recently developed techniques such as the use of genetically engineered bacteriophage to display and/or secrete the χ-protein and thereby select for an improved confirmation. (Marks, J.D., et al., J. Mol. Biol. 222:581-597 (1991)).

Additional optimizing techniques, as described above, may also be used to confer "special" properties on the χ -protein, such as "humanizing" the χ -protein 15 using human FR-like regions as the scaffold protein to reduce the possibility of adverse immunologic reactions during therapy. (Daugherty, B. L., et al., Nucleic Acids Res. 19:2471-2476 (1991)). Special properties may also include increased solubility or stability or improved renaturability or secretion.

This invention further relates to the method of producing the χ -protein which includes the steps of determining the splice points for additional binding sites, as described in Examples 2 and 3, determining the amino acid sequence of the resulting construct, deducing the DNA sequence encoding that amino acid sequence, inserting the DNA sequence into an appropriate expression vector and expressing the protein in a suitable host system, refolding the protein to its biologically active conformation and analyzing its biological binding activity. For example, in the case of a χ -protein with catalytic activity, this would include measurement of enzymatic properties.

-27-

The ability to target therapeutic agents in a host with antibodies has been a long-term goal of medical research. The term host, as used hereinafter, is intended to encompass mammalian hosts, including humans. The most elegant targetable proteins would consist of the minimum structures needed for selective delivery and effector function. (Tai, M., et al., Biochem. 29: 8024-8025 (1990)). Although the utility of binding proteins having multiple binding sites of different specificity has been widely recognized (U.S. Patent 4,676,980), these crosslinked antibodies or crosslinked antibody fragments suffer from the need for complex purification schemes and large molecular size.

5

10

35

Notwithstanding the construction of the smallersized sFv, the need still exists for a multifunctional 15 binding protein with as reduced a size and as compact a shape as possible to permit effective therapeutic use. The subject invention also relates to use of these χ proteins in therapeutic and diagnostic procedures. These uses include, but are not limited to, in vivo and 20 in vitro imaging agents, delivery agents for drugs, radioisotopes, and cytotoxic substances. The χ-protein could also include a binding site for an effector molecule, such as an enzyme, growth factor, celldifferentiation factor, lymphokine, cytokine, hormone, 25 anti-metabolitic, or an ion-sequestering sequence such as calmodulin. The χ -protein could further include a binding site reactive with antibody-dependent cytolytic cells or cytotoxic T cells. Also included are χ -proteins with catalytic or biosensor activity, as 30 well as binding sites that facilitate affinity purification procedures.

In some instances, the primary binding site could be a high affinity site that targets the protein to specific cell surface locations, and the secondary site

15

20

25

30

35

designed to decrease normal receptor activity. A χ -protein could be designed with one binding site comprising a receptor for a cell surface antigen and a second binding site comprised of a modified receptor with diminished affinity for its ligand. Thus the χ -protein would to bind a cell via the normal binding site, leaving the binding site with diminished binding activity exposed, thereby resulting in decreased receptor activity. For example, a χ -protein could be designed with neural cell adhesion molecule (NCAM) variable regions that would modulate cell interactions such as contact inhibition of malignant cells.

A χ-protein may also be designed to modify, enhance or inhibit cell-cell interactions, or communication. For example, one binding site could be designed to target a cancer cell and a second binding site could be designed to target an effector cell, such as a macrophage, thus binding the macrophage in a manner which results in destruction of the targeted cell.

Moreover, a χ -protein could mediate phenomena such as antibody-dependent cellular toxicity. (Huston, J.S., et al., Proc. Natl. Acad. Sci. USA, 85:5879-5883 (1988)). For example, a χ -protein could be designed with one binding site comprised of a receptor for a cell surface marker protein and a second site derived from a receptor for killer T cells. The χ -protein would bind to the target cell and the secondary site would bind the cytotoxic T cell, resulting in destruction of the target cell. The χ -proteins of the present invention would exhibit significantly greater tissue accessibility or tumor penetration and faster pharmacokinetics due to their compact size.

Recently, it has been shown that an Fv undergoes a conformational change upon antigen binding. (Bhat, T.

-29-

N., Nature, 347:483 (1990)). It is reasonable to predict, that the χ -proteins of the present invention would undergo similar conformational changes, due to the significant sequence and structural homology with the native Fv. Although this conformational change is modest, it involves the translational movement of the V_H and V_L domains relative to each other, such that the orientation of the bottom loops can shift. This makes the χ -protein conformation sensitive to binding at either the first or second binding site, and requires that linkage exists between their binding equilibria. It is reasonable to predict that this conformational change would enable the χ -protein to act as a "molecular switch".

5

10

15

20

25

30

For example, the first site of the χ -protein may be a catalytic antibody combining site (e.g., a site that catalyzes the conversion of a "pro-drug" to a cytotoxic drug), while a second site is specific for binding to a marker on a cell surface (e.g., a tumor cell). If binding to the cell surface epitope induces a conformational change in the χ -protein such that the loops of the catalytic site are brought into optimal geometry for efficient catalysis, then the pro-drug would be converted to cytotoxic drug directly at the site of the tumor, so that high cytotoxic levels could be maintained at the site while serum levels remained low. The unbound catalyst would have low efficiency and, because of its low molecular weight, would clear rapidly from circulation.

The χ -proteins are well suited to applications in immunotargeting because of their binding capabilities and compact size. The absence of constant domains will reduce nonspecific background binding, thus enhancing visualation of target tissue by <u>in vivo</u> or <u>in vitro</u>

10

imaging procedures. The use of χ -proteins in <u>in vitro</u> diagnostics would also reduce nonspecific binding thereby increasing the accuracy of these assays.

The χ -proteins of the present invention are also useful to deliver drugs, cytotoxic substances or effector molecules to immunotargeted tissues. One of the binding sites of the subject protein may exhibit catalytic activity that is triggered by binding of a specific ligand to the other binding site.

The invention will be further illustrated by the following Examples, which are not intended to be limited in anyway.

Example 1

Constructing a Model of Bivalent McPC603 xsFv

The following illustration, which utilizes the 15 McPC 603 Fv region (SEQ ID NOS: 1 and 10), for which the three-dimensional crystallographic structure has been determined, has been chosen in order to assess how well a particular set of splice points will generate a χ -site (additional binding site) that recreates the parent binding site. This example also reveals the general method of building a χ -protein model, by the specific example of mapping the McPC603 $V_{\rm H}$ CDRs to $V_{\rm H}$ bottom loop positions, and the V_L CDRs to V_L bottom loop positions. Molecular modeling was performed on a 25 Silicon Graphics 4D/70GT Superworkstation using the Biosym programs INSIGHT (to visualize models), HOMOLOGY (to assemble spliced CDR/FR sequences), and DISCOVER (to minimize the energy of the model) (Biosym, Inc., San Diego, CA). The HOMOLOGY program was used to 30 assemble the structure in five steps, and the DISCOVER program was used to minimize the energy of the resulting model about the splice points. Although,

-31-

HOMOLOGY was used in this example, this program is not necessary to build the model. Any one of several alternative molecular mechanics programs, such as AMBER and CHARMM, could be used instead of the DISCOVER program.

5

10

Two copies of the McPC603 sFv structure were superimposed (Figure 3A): model A had the binding site up and the model B had the bottom loops up; the CDR loops of B superimposed on the lower loops of A, as described above. The primed loops (e.g. H2') are those spliced into the χ -site.

Building the H2' loop and the bridge. (Figures 4A 1. The A coordinates for the $V_{\rm H}$ domain were and 4B). used up to where H2' from structure B splices into the C-terminus of structure A and the A 15 coordinates of the linker and the following Vi domain were likewise used. The coordinates of the H2' segment were taken from B. Once coordinates had been assigned to the residues flanking the bridge peptide region, the HOMOLOGY loop search 20 algorithm was used to find a 5 residue peptide whose flanking 3 alpha carbons overlapped well with the 3 amino acids on either side of the gap, and whose conformation best fit the surface of the V, domain. The HOMOLOGY program automatically 25 connects the peptide bonds at the splice points to create the new single chain structure called MDL1. It will generally be necessary to rotate side groups in the interface between VH and H2' out of the way in order to reduce steric conflicts. One 30 residue in H2' had to be changed because simple rotation could not eliminate its steric conflicts: Phe 137 (position H70)-> Ala. Using the DISCOVER

program, the atoms about the splice points were moved so as to minimize contributions to the energy of the structure due to improper bond lengths and geometries. At this point the composition of bridge peptide was polyalanine; additional computational analysis of the model may be used to further assess the optimum length, path and composition of the bridge. This new structure (with H2' and the bridge) is called MDL1.

- 2. Building H3' and H1' Loops. (Figures 4C and 4D)
 Structure MDL1 supplied the coordinates for the
 residues preceding and following the segment where
 H3' was to be built into the model. The
 coordinates for the H3' segment were taken from
 structure B. Since H1' was a short segment of 3
 residues that would replace exactly 3 residues of
 MDL1, H1' was built by side chain substitution.
 This structure is MDL2.
- 3. Building L3'. (Figure 4E) Structure MDL2 supplied the coordinates for the residues preceding and following the segment where L3' was to be built into the model. The segment for L3' was built into the next stage of the model in a manner analogous to the way that MDL2 was built. The resulting structure is MDL3.
- 4. Building L1'. (Figure 4F) Structure MDL3 supplied the coordinates for the residues preceding and following the segment where L1' was to be built into the model. The segment for L1' was built into the next stage of the model in a manner analogous to the way that MDL2 was built. This structure is MDL4.

-33-

5. Building L2'. (Figure 4G) Structure MDL4 supplied the coordinates for residues preceding the segment where L2' was to be built into the C-terminus of the model. The segment for L2' was built into the next stage of the model in a manner similar the way that MDL1 was built except that no connection (bridge or linker) was made to the C-terminus of the L2' segment. In order to stabilize the C-terminus of the L2' loop, a disulfide bridge was constructed between the C-terminus of L2' and the segment that derived from the V_L N-terminal strand by substituting the Arg in L2' (at the position corresponding to L68 in V_L) and the Thr at the position corresponding to L5 in V_L with Cys.

5

10

Refinement. At each stage, atoms making bad 15 contacts were rotated out of the way. This often required breaking the peptide bond between neighboring residues, rotating a backbone bond and then reforming the peptide bond. functionality for these modeling operations is 20 built into the INSIGHT program and other similar molecular modeling programs. The resulting modified peptide bond was added to the list of splice points. Strain in the splice point peptide bonds was minimized by subjecting the residues on 25 either side of the spliced peptide bond to 100 cycles of steepest descent minimization using the DISCOVER molecular mechanics program. The final structure was minimized for 1000 steps of steepest descent minimization without coulumbic 30 interactions, followed by 200 steps steepest descent, and 1000 steps of conjugate gradient

minimization with columbic charge interactions.

15

30

Figures 4A-4G show the final model of the McPC603 (McPC603) $V_H(V_H) - V_L(V_L)$ χ -protein with corresponding pairs of loops highlighted with ribbons. The symmetry between the top and the bottom of the molecule can be seen. Each of the 5 pairs of spliced loops are highlighted in separate frames.

A detailed comparison of the conformation of the new sites is presented in Figure 5. The upper view is of a first combining site on the top of the molecule while the lower view is the χ -site built onto the bottom of the molecule. In the figure, corresponding regions of the loops are highlighted and side chain heavy atoms are shown. It can be seen that during the minimization process, there was some divergence in the conformation of the backbone and of the side chains between these two sites, but there still remains an impressive degree of homology between them. This is a measure of the congruency of the lower loop stems and the corresponding upper loop stems. The observed homology also suggests that the steric environment surrounding the spliced loops must be similar to that of the parent loops, even though it is clear that the loops of the first site are more exposed than are those of the second x-site.

25 Example 2 Defining top and bottom loop symmetry and choosing splice points.

Because of the high level of symmetry that exists between the V_H and V_L subunits, CDR loops can be spliced in a "homologous" fashion (V_H top loops onto V_H bottom loops, $V_H(V_H)$, and V_L top loops onto V_L bottom loops, $V_L(V_L)$) or in a "heterologous" fashion (V_L top loops onto the bottom of V_H , $V_H(V_L)$, and V_H top loops

onto the bottom of V_L , $V_L(V_H)$). In addition, there are two ways to connect the V_H and V_L regions: 1) linker peptide bridging from the C-terminus of V_H to the N-terminus of V_L (V_H - linker - V_L), or 2) linker peptide running from the C-terminus of V_L to the N-terminus of V_H (V_L - linker - V_H). There are, therefore, four possible types of χ -protein derived from immunoglobulin CDRs.

If one is splicing the CDRs of one V_H into the bottom loops of another V_H (as in Example 1) then H2 would be spliced into the V_H C-terminal strand. The C-terminal end of the new H2' segment would end up on the outside of V_H region, near its N-terminal strand (Figure 3). Thus, if one is splicing heavy chain loops into the bottom of the heavy chain in a sFv connected in the V_H - V_L order, the C-terminal end of H2' can be connected to the peptide that links the heavy and light chains. The L2 loop structure can likewise be spliced onto the C-terminus of V_L .

10

15

25

30

Modeling a $V_H(V_H) - V_L(V_L) \chi$ -Protein based on McPC603:

The following discussion presents a detailed analysis of how to construct a second McPC603 antigen binding site on the bottom of McPC603 to form a homodomain V_H -linker- V_L χ -protein. Although this exemplification is based on McPC603, the structural homology that exists between the framework regions of known Fv structures clearly suggests that CDR splicing to the bottom loops of Fv regions will work with any antibody Fv. The existing structural homology between known structures permits one to structurally align the sequences of these known structures and to establish a numbering index wherein the position numbers in the strand regions identify unique structural locations.

The position scale used to refer to splice points is defined in Figure 2B.

Determining loop splice points

When the CDR loops are spliced into the lower loops, it is important to preserve those framework residues bordering both the top loops and the bottom loops which are critical for maintaining the proper folding of the β-barrel structure. In the following figures, the spatial alignment of bottom loops with the superimposed CDR loops are shown along with the segment of aligned sequence that derives from the chosen splice points. Three considerations govern the choice of the splice points:

- Preservation of critical framework residues.
- 15 2. Maximal incorporation of CDR sequences.

-37-

3. Fusion at closet possible splice points. In order for the spliced loops to maintain their native structure in the new site, it is important that the chains be distorted as little as possible at the splice sites. Therefore one tries to chose splice points as close as possible to the positions where the aligned chains are closest together, e.g., where corresponding alpha carbons are separated by no more than a few angstroms.

Determining the Splice Points for Splicing H3 (H-TL4) into H-BL2, and L3 (L-TL4) into L-BL2:

5

15

20

25

30

Figures 6 and 7 show how the CDR 3 regions (H3 and L3) would be spliced into H-BL2 and L-BL2, respectively. Critical framework residues H-BL2 and L-BL2 include the Gln at position H39 in H-BL2 and the Gln at position L45 in L-BL2.

Referring to Figure 6, residues position H102 to H114 from H3 (positions H101 to H115) can be spliced into H-BL2. The H3 residue at position H101 is not included because it would replace the critical framework residue at position 39. The H3 residue at position H115 is not included because splicing from position H114 in the H3 loop to position H47 in the framework leads to less distortion of both the framework and the loop conformations.

The L3 loop (positions L97 to L104) can be spliced into L-BL2 in a similar manner. Referring to Figure 7, the new L3' loop includes residues from positions L99 to L104. The L3 residues at position L97 to L98 are not included because L98 would replace the critical framework residue at position L45. The two chains are sufficiently close together at L3 position L104 that the entire C-terminal end of L3 can be spliced into the new L3' loop. Consequently, only one residue from the

10

25

30

N-terminal end of L3 cannot be spliced. In the cases of both the H3' and the L3' splices, the backbones of the respectively aligned chains lie close enough together at the indicated splice points that there is little distortion of the backbone conformation in the χ -protein.

Determining the Splice Points for Splicing H1 (H-TL1) into H-BL4, and L1 (L-TL1) into L-BL4:

Figures 8 and 9 show how the CDR1 regions would be spliced into the fourth bottom loops. Because sections of both H-BL4 and L-BL4 that contain non-critical residues do not align well with the CDR1 portion of either CDR loop, only portions of the CDR1 loops can be spliced.

The H1 segment (positions H31 to H35) is 5
residues long in McPC603, and of these, the 3 residues
(positions H31 to H33) that can be spliced are on the
outer portion of H-TL1, which faces the binding site.
Position H92 in H-BL4 is a critical residue that is
involved in a conserved intrachain salt bridge, while
positions H96 to H98 are conserved inner strand
residues. Consequently positions H31 to H33 are
spliced between positions H92 and H96 in H-BL4 to
create H1'.

In McPC603, L1 (positions L25 to L41) is fairly long (17 residues), but only the internal 10 residues (position L30 to L39) can be spliced (Figure 9).

Position L30 is the first place where the L1 loop and L-BL4 align. Framework positions L93 to L95 are conserved, inner strand residues that are structurally analogous to H96 to H98. In order to preserve framework position L93 the L1 loop is spliced at position L39. Consequently, 10 residues from the McPC603 L1 region can be spliced into L-BL4. In other

antibodies having shorter L1 regions, only about 3 or 4 residues can be spliced.

<u>Determining the Splice Points for Splicing H2 (H-TL2)</u> into the C-terminal end of VH:

5

10

15

20

25

30

Figure 10 shows how the H2 region of the inverted copy of McPC603 aligns with the N-terminal and the Cterminal ends of V_H. Residues positions H48 to H50 that flank the N-terminal end of H2 loop region (positions H50 to H70) align closely with V_{H} C-terminal residue positions H117 to H119, respectively. When splicing in the H2' loop, one wants to preserve as much of the $V_{\rm H}$ C-terminus as possible. Based on the superposition of the C-terminal and H2' backbones, splicing can be done between positions H115 and H119 in the C-terminal strand and between positions H47 and H51 in N-terminal flanking sequence of H2. Because several residues in the C-terminal sequence of V_{H} (and V_{L} as well) are probably important to stabilizing the packing of BL1 and BL4, the splice should be made so as to preserve as much of the C-terminus as possible. As shown in Figure 10, therefore, the splice is made between position H119 of the V_H C-terminal strand and position H51 of the H2' loop. The C-terminal end of H2' (arrow A, Figure 10) is located on the surface of V_H , near the N-terminal strand of V_H , when H2' is properly positioned relative to H3', L3', H1' and L1'. To connect this end of the N-terminal end of the interchain linker used to create an single chain Fv (arrow B, Figure 10), a 5 residue bridge peptide must be added (between points B and A in Figure 10). In this arrangement, the linker-bridge structure folds across the bottom of the $V_{\rm H}$ chain. The preferred bridge peptide would contain Ser for solubility and non-polar

10

15

20

30

residues to help anchor it to the surface of $V_{\rm H}$ so as to stabilize the conformation of the H2' fold. In the model and in the sequence shown in Figure 12, a polyalanine bridge has been used.

Figure 11 shows how the region of the inverted copy of McPC603 aligns with the N-terminal and Cterminal ends of V_L . Residue positions L49 to L57 that flank the L2 loop region (positions L57 to L63) align closely with V_L C-terminal residue positions L102 to L109. The same considerations are involved in making the L2' splice as with the H2' splice. The splice is made from C-terminal position L108 to L2 position L56. The L2 loop structure folds around until it runs parallel to the V_L N-terminal strand. In order to stabilize the L2' loop structure, we include a disulfide bond from the end of the L2' loop to a residue in the N-terminal strand of V_L . To accomplish this the Arg at position L68 in the C-terminal end of the L2 loop and the Thr at position L5 in the Nterminal strand of V_L were changed to Cys.

Figure 12 shows the sequence (SEQ ID NO: 17) of the resulting construct with the sequences of the primary CDR sites (bold) and secondary CDR sites (Underlined) identified.

25 Example 3 Determining χ-Site Splice Points From Primary Sequence Alignment

The members of the Ig Superfamily share a high degree of structural homology. By using the homology that exists between members of known structure, it is possible to engineer χ -proteins from Ig Superfamily molecules whose sequence is known but whose tertiary

-41-

structure is not yet determined by crystallographic or NMR methods.

Figure 13A (SEQ ID NOS: 10-16 and 18-30) and 13B (SEQ ID NOS: 1-9 and 31-40) shows the set of sequences of Fv domains for which structures are known and are available from the Brookhaven Protein Databank (PDB at Brookhaven National Laboratory).

5

Table 1 lists the sequence and structure identification names, along with the Kabat et al. classification and references for the sequences that 10 appear in Figures 2B (SEQ ID NOS: 1-16) and 13A (SEQ ID NOS: 10-16 and 18-30) and 13B (SEQ ID NOS: 1-9 and 31-The alignment of these sequences is based on the structural homology that exists between them, 15 especially in the β -strand regions (OS and IS). Because of this high degree of structural correlation, one can assign positions in a "consensus 3-D model" to residues in a set of aligned β -barrel sequences. Furthermore, at certain positions, there exists a 20 strong preference for certain residues or for residues having certain physical characteristics (Table 2). Thus, while the 3-D structures of most Fv sequences remain undetermined experimentally, the residue preferences at these positions makes it possible to 25 reliably align these sequences with those V or Ig Superfamily sequences for which the 3-D structures have been experimentally determined. Ig V regions can thereby be aligned with the sequences of more distant members of the immunoglobulin superfamily, such as the 30 T-cell receptor which was successfully modeled according to the McPC603 Fv structure (Novotny, J. et al. Proc. Natl. Acad. Sci USA 88:8646-8650 (1991).

Specificity	PDB ¹ ID	Kabat² Id	KABA Ligh page	t Cha	KABAT Light Chain Ref. page ∉ Group	KABAT Heavy Page	r P B Cha	KABAT Heavy Chain Ref. page # Subgroup	SOURCE
Ant1-Phoshorylcholine	тись	MCPC603 ^{3,4}	151	151 . 23	KAPPA I	431	28	III(A)	Mouse
Anti-Hen Egg White Lysozyme	1FDL ⁵	D1.3¢	214	.133	214 133 KAPPA V	363	8	I(B)	Mouse
Anti-P-Azobenzenearsonate Hybri.	15197	R19.97	1282	36	KAPPA V	1443	43	V(A)	Mouse
Anti-beta 1,6 D-Galactan	2FBJ ⁸	J539°	244	129	KAPPA VI	452	23	III(A)	Mouse
Anti-Fluorescein Hybrid.	4FAB10	4-4-2011	165	47	KAPPA II	459	27	III(C)	Mouse
Anti-Hen Egg White Lysozyme	2HFL ¹²	HYHEL-5 ¹³	241	63	KAPPA VI	500	ີຕ	MISC.	Mouse
Anti-Hen Egg White Lysozyme	JHFK16	HYHEL-10 ¹³	216	182	KAPPA V	340	21	I(A)	Mouse
(B-J Fragment)	1REI 15	REI ¹⁵	104	38	KAPPA I	:	:	:	Нитал
(B-J Fragment)	2 RHE 14	RHE ¹⁶	131	23	LAMBDA I	•	:	:	Human
(IGG1-Lambda)	2FB417	KOL ^{17,18}	131	17.	LAMBDA I	326	53	III	Нишал
Anti-Digoxin Hybri.		26-1019,20	165	£3	KAPPA II	383	188	II(A)	Mouse
Anti-Lysozyme G-loop Hybri.	:	GL00P421.	:	:	•	:	:	:	Mouse

-43-

Table 1 Notes

- Bernstein, F.C., Koetzle, T.F., Williams, G.J.B., Meyer, E.F., Jr., Brice, M.D., Rodgers, J.R., Kennard, O., Shimanouchi, T., & Tasumi, M. (1977). J. Mol. Biol., 112, 535-542.
- E.A. Kabat, Tai, T.W., Perry, H.M., Gottesman, K.S. and Foeller, C. (1991). "Sequences of Proteins of Immunological Interest", Vols. I-III. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, Publication No. 91-32421.

5

20 .

- Rudikoff, S., Satow, Y., Padlan, E., Davies, D., &
 Potter, M. (1981). <u>Mol. Immunol.</u>, 18, 705-711.
- Potter, M., Rudikoff, S., Padlan, E. & Vrana, M.
 (1977). In "Antibodies in Human Diagnostics and Therapy", Raven Press, New York.
 - Fischmann, T.O., Bentley, G.A., Bhat, T.N.,
 Boulot, G., Mariuzza, R.A., Phillips, S.E.V.,
 Tello, D. & Poljak, R.J. (1991). J. Biol. Chem.,
 266, 12915 12920.
 - Verhoeyen, M., Milstein, C. & Winter, G. (1988).
 Science, 239, 1536-1538.
- Lascombe, M.-B., Alzari, P.M., Boulot, G., Saludjian, P., Tougard, P., Berek, C., Haba, S., Rosen, E.M., Nisonoff, A., & Poljak, R.J. (1989). Proc. Nat. Acad. Sci. USA, 86, 607-611.

- 8. Suh, S.W., Bhat, T.N., Navia, M.A., Cohen, G.H., Rao, D.N., Rudikoff, S. & Davies, D.R. (1986).

 Proteins, Structure, Funct., 1, 1986.
- 9. Rudikoff, S., Rao, D.N., Glaudemans, C.P.J., & Potter, M. (1980). Proc. Nat. Acad. Sci. USA, 77, 4270-4274.
 - 10. Herron, J.N., He, X., Mason, M.L., Voss, E.W. Jr., & Edmundson, A.B. (1989). Proteins Struct.

 Funct., 5, 271-280.
- Bedzyk, W.D., Johnson, L.S., Riordan, G.S. & Voss,
 E.W., Jr. (1989). J. Biol. Chem., 264, 1565-1569.
- Sheriff, S., Silverton, E.W., Psdlan, E.A., Cohen, G.H., Smith-Gill, S.J., Finzel, B.C. & Davies, D.R. (1987). Proc. Nat. Acad. Sci. USA, 84, 8075-8079.
 - Smith-Gill, S.J., Mainhart, C.R., Lavoie, T.B.,
 Rudikoff, S. & Potter, M. (1984). <u>J. Immunol.</u>,
 132, 963-967.
- 14. Padlan, E.A., Silverton, E.W., Sheriff, S., Cohen,
 20 G.H., Smith-Gill, S.J., & Davies, D.R. (1989).
 Proc. Nat. Acad. Sci. USA, 86, 5938-5942.
 - Palm, W. & Hilschmann, N. (1973) Z. Physiol.
 Chem., 354, 1651-1654; (1975) Z. Physiol. Chem.,
 356, 167-191.
- 25 16. Furey, W. Jr., Wang, B.C., Yoo, C.S. & Sax, M. (1983). <u>J. Mol. Biol.</u>, 167, 661-692.

PCT/US93/04338

10

15

- 17. Kratzin, H.D., Palm, W., Strangel, M., Schmidt, W.E., Friedrich, J. & HilschMann, N. (1989).

 Biol. Chem. Hoppe-Seyler, 370, 263-272.
- Schmidt, W.E., Jung, H.-D., Palm, W. & Hilschmann,
 N. (1983). Z. Physiol. Chem., 364, 713-747.

 - 20. Mudgett-Hunter, M., Anderson, W., Haber, E. & Margolies, M.N. (1985). Molec. Immun., 22(4), 477-488.
 - 21. Darsley, M.I. & Rees, A.R. (1985). <u>EMBO J.</u>, 4(2), 393-398.

Once a sequence has been positioned in relation to the set of structurally aligned V region sequences, one can use the splice point analysis developed in Example 2 to construct a model and delineate a sequence for a χ -protein by analogy. The procedure for determining proper slice points is as follows:

Sequences of undetermined 3-D structure are lined up with the set of structurally aligned sequences. The sequence alignment process makes use of the conserved and semi-conserved residue positions listed in Table 1 to fix the alignment in certain regions. Between these anchor points it may be necessary to introduce gaps. Sequences in the β-strand regions (OS and IS) generally line up without the need for introducing gaps except in the N-terminal section of the light chain variable region (L OS1 - L BL1). Relative to all other

PCT/US93/04338

5

10

15

20

25

30

light chain V region classes, kappa light chain class V sequences have a deletion at position L7 and an insertion at position L17. Lambda light chain sequences also have a deletion at position L7, but no insertion at position L17. Kappa light chain class VII sequences have a deletion at position L23. Other than these gaps in the Nterminal section of the light chains, the alignment of sequences is may make use of conserved positions to establish anchor points. Further help in the alignment of intervening sequences is derived from conservation of sidechain properties (i.e., polar, non-polar, charged, acid, basic, etc) at certain positions. The other major gaps will exist in the CDR loop regions (H1, L1, H2, L2, H3 and L3). Because CDR loops are variable in sequence and length, the exact location of gaps in these regions is less critical as the sequences in these regions show little homology. Structural variation tends to be the largest at the center of CDR loops of similar sequence, so gaps are preferably positioned in the center of CDR top loops.

2. Once the sequences of one or more V regions of undetermined structure have been lined up with the structurally aligned reference set, structure positions are assignable to the residues that locate in the regions of the inner and outer β -strands. Because the χ -site splice points occur near the ends of the β -strand regions, they can be assigned by analogy with the splice positions determined for McPC 603 in Example 2. Given two sequences aligned with the reference set, the primary sequence is that into whose bottom loops

-47-

the χ -site will be built, and is referred to below as the target sequence; while the secondary sequence contains the CDR loops which will be built into the χ -site, and which will be referred to below as the source sequence. 'x-site loops are created by splicing top loop segments from the source sequence in place of bottom loop segments in the target sequence. For a $V_H(V_H) - V_L(V_L) \chi$ protein construct similar to that designed in Example 2, the spliceable portions of source CDRs, flanked by the fixed top loop β -strand end points, are labelled in Figure 13 as L1'(S), L2'(S), L3(S), H1'(S) and H3'(S), while the target bottom loop segments, flanked by the fixed bottom loop β strand end points, are labelled in Figure 13 as L1'(T), L2'(T), L3'(T), H1'(T), H2'(T) and H3'(T). Thus, to create a χ -site L3' loop in the L BL2 loop of the target sequence, one would splice the residues identified as L3'(S) in Figure 13 to the residues flanking the segment identified as L3'(T); similarly, to create a χ -site H1' loop in the H BL4 loop of the target sequence, one would splice the residues identified as H3'(S) in Figure 13 to the residues flanking the segment identified as H1'(T), and so forth. These splices are done to create a $V_H(V_H) \cdot V_L(V_L)$ χ -protein as in Example To create a $V_H(V_L) \cdot V_L(V_H)$ χ -protein L3' loops in the H BL2 loop of the target sequence, one would splice the residues in the segment identified as L3'(S) in Figure 13 to the residues flanking the segment identified as H3'(T). Likewise, to create a χ -site H1' loop in the L BL4 loop of the target sequence, one would splice the residues flanking the region identified as H1'(S)

30

5

10

15

20

10

15

20

25

30

in Figure 13 to the residues flanking the segment identified as L1'(T), and so forth.

Alternatively, one can build structural 3. models for the sequences of undetermined structure using the coordinates of one or more of the known structures as a basis for the framework residue locations. Programs such as the Biosym HOMOLOGY (Biosym, Inc., San Diego, CA) program are designed to aid in this process. The procedure is well established in the literature (Grear, J. (1991) Meth. in Enzymol. 202:239-252 (1991)). Once constructed, models of the parent Fv, sFv or V region and of the Fv regions of the corresponding x-site can thus be used to determine splice points and to construct a model of the χ -protein in the manner set forth in Examples 1 and 2. Building a model provides insight into possible steric conflicts, particularly, as was noted in Example 2, at the interface between H2' and the surface residues of the Fv domain.

Figure 14 shows the sequence of χ -protein R19.1 (D1.3) (SEQ ID NO: 42) as it would be constructed following parts 1 and 2 of the example using the aligned sequences in Figure 14. The framework and primary CDR loops are taken from R19.9 while the sequences for the χ -site loops were taken from the H1'(S), H2'(S), H3'(S), L1'(S), and L3'(S) regions of D1.3.

Figure 14 also shows the sequences of χ -protein MCP(MCP) (McPC603 (McPC603) (SEQ ID NO: 41)) as it would be constructed in Examples I and 2. Also shown are sequences of χ -proteins 26-10(D1.3) (SEQ ID NO: 43) and

-49-

26-10(GL00P4) (SEQ ID NO: 44) as they would be constructed using the method of Example 3.

TABLE 2

Light Chain Position	Residue Preference	Heavy Chain Position	Residue Preference
L1	D, (E)	Н1	E, (D,Q)
L2	I	H2	v
		нз	Q,K,H
L4	M,L	H4	L
L5	т		
L6	Q	Н6	E(Q)
L7	S,T,(-)	H7	S, (T,P,-)
L8	P, (T,E)		
		H11	L, (V)
	•	H12	V, (M)
		H14	P, (A)
L16	G(L)	H15	G,S
L17	-,G		
L20	V, (A)	H18	L,V,(I)
L21	T(S)	H19_	K,R,S
L22	I, (M,L)	H20	L,I,M
L23	S,T,(-)	H21	S,T
L24	С	H22	С
L27	S, (T)		
		H25	S,T
		H26	G
		H27	Y,F,(D,T,-)
		H28	polar
		H29	F,I,L
		Н30	T,S,(D)

-50-

TABLE 2 CONT

]
Light Chain Position	Residue Preference	Heavy Chain Position	Residue Preference
L40	L,M,V,(A)		,
L42	W	Н36	W
L43	Y, (V, I)	Н37	·V,(I,M)
L44	Q	Н38	K,R
L45	Q, (K)	Н39	Q, (K)
L46	K		·
L47	P,(S)	H41	P, (H)
	·	H42	G, (E)
		H43	K,Q,N,R
L51	P, (V,F)	H45	L
L52 ·	K, (R,Q)	H46	E, (D)
		H47	W, (Y,H,D)
		H48	I, (M,L)
L54	L,(W)		
L55	I,(V)	H49	G, (A)
L56	Y, (G,K)		
		H51	I,(V,S)
L59	s		
		H62	Y, (T)
L64	G		
L65	V, (I)	Н66	F,(L,V)
L66-	P, (S)	Н67	K, (Q,M)
L68	R	Н69	K,R,(L)
L69	F		
L70	S, (T)		

-51-TABLE 2 CONT

	T		
Light Chain	Residue Preference	Heavy Chain	Residue Preference
Position	reference	Position	·
L71	G, (A,V)	H72	F,I,L,V
L72	S	Н73	S,T
L74	s	H75	D, (N)
L75	G		
L76	T	н77	S, (T, A, P)
<u> </u>		Н78	polar
		Н79	S,N
		H80	(-)
· ·	:	H81	A,L,V
		H82	Y, (F,H)
L80	L, (F)	Н83	L, (M)
		Н84	Q, (D, E, K)
L82	I	H85	L,M,I
L83	polar	H86	S,N, (D,R)
L85	V,M,L,(A)	Н88	L,(V)
L86	Q,E		
L88	E	Н91	E, (D, A)
L89	D	H92	D
		н93	T,S
L91	A, (G)	H94	A, (G)
L93	Y, (H)	Н96	Y
L94	Y,F	Н97	Y, F
L95	С	Н98	С
L96	Q, (A,S)	Н99	A
	•	H114	D,A

-52-

TABLE 2 CONT

Light Chain Position	Residue Preference	Heavy Chain Position	Residue Preference
L106	T, (E, V)		,
L107	F, (P)	H116	W
L108	G	H117	G
·		H118	Q,A
L110	G	H119	G
L111	T	H120	Т
L112	K		
L113	L, (V)	H122	V, (L)
L114	E, (T,Q,K)	H123	T .
L115	I, (L,V)	H124	V
L116	K	H125	s

Notes:

5

- 1) Single letter amino acid codes; letters in parentheses signify less common preferences.
- 2) "-" signifies the occurrence of a gap.
- 3) Each row corresponds to an equivalent location in the V_H and V_L chains.

EXAMPLE 4

Construction, Expression and Evaluation of a χ -Protein

The χ -1 and χ -2 genes (Figure 15A and B, SEQ ID NOS:
45 and 47) were prepared by mutagenesis of the 26-10 sFv

10 gene as described in (Huston, J.S., et al., Proc. Natl.

Acad. Sci. USA, 85:5879-5883 (1988); Tai, M.-S., et al.,

Biochemistry 29:8024-8030 (1990)) and were incorporated into the pET vector described in Studier, F.W., and Moffat,

B.A., J. Mol. Biol. 189:113 (1986) behind a T7 promoter.

15 Upon transformation of E. coli with this vector, direct

expression produced each in the form of cytoplasmic inclusion bodies. Cells were treated with lysozyme to allow cell lysis and ultracentrifugal isolation of the inclusion bodies, which were then dissolved in 6 M 5 quanidinium chloride containing 10 mM dithiothreitol, 25 mM Tris, and 10 mM EDTA at pH 8.1; the solution was incubated overnight at room temperature. The protein was then diluted into 3 M or 3.5 M urea buffer containing 25 mM Tris, and 10 mM EDTA, and a glutathione redox couple (1 mM 10 oxidized, 0.1 mM reduced) at pH 8.1. Following 18 h. at 4°C, each was fully renatured by dialysis into phosphate buffered saline (PBS), consisting of 0.05 M potassium phosphate, 0.15 M NaCl, pH 7.0, and 0.03% NaN3. and χ -2 solutions were then passed through ouabain-15 Sepharose columns, washed first with PBSA (PBS +0.03% NaN3), then with 1M NaCl in PBSA to remove any unbound material from the columns, and elution effected by displacing specifically bound protein with 20 mM ouabain in The affinity purified χ -1 and χ -2 proteins were then 20 examined by SDS polyacrylamide gel electrophoresis. Figure 16A and B shows the χ -1 (SEQ ID NO: 46) and χ -2 (SEQ ID NO: 48) polypeptide chains following their affinity isolation. In figure 16B, the oxidized χ -2 (upper band) is compared with 26-10 sFv (lower band) in lanes denoted as mixture. 25 Sequence analysis of the χ -2 protein verified that the insertions noted in the gene sequence (Figure 15B) were present in the protein sequence CNBr fragments of the protein were made and sequenced as a mixture in an Applied Biosystems 470A gas-phase sequencer equipped with a model 30 120A on-line analyzer. The gel (Figure 16B) is also consistent with the increased molecular weight of the χ -2 over the 26-10 sFv. Refolded protein that bound to the

ouabain-Sepharose column has necessarily regained active

antibody combining sites for digoxin-like cardiac glycosides typified by ouabain.

Additional insights into the shape and properties of the χ -1 and χ -2 proteins are apparent from Superdex 75 size 5 exclusion chromatography of the affinity purified χ proteins in comparison to the 26-10 sFv, as shown in Figure 17. The single H1' insertion of χ -1 adds two tyrosyl residues within the GYGY sequence, resulting in a pronounced tendency to dimerize and a very skewed profile 10 indicative of dissociation into monomer (data not shown). In contrast, data shown in Figure 17 for χ -2 (top panel) indicate that it appears perfectly behaved in solution, devoid of any apparent dimer. Thus, as one incorporates a multiplicity of χ -CDRs in V regions, the known 15 hydrophobicity of CDR sequences apparently is contained by the aggregate of χ -CDR conformation and interaction. added H1' and H2' loops necessarily increase the protein's Stokes radius, resulting in its elution position being between the 26-10 monomer and dimer positions (bottom 20 panel). A mixing experiment that combines both proteins in a single chromatographic separation (middle panel) indicates that the χ -2 shows no apparent interaction with the 26-10 monomer and dimer species, as the middle profile appears to be a simple additive composite of the top and 25 bottom chromatograms. The peaks beyond 30 minutes (horizontal axis is in minutes) are simply injection artifacts on the HPLC system, probably due to buffer differences between the column and sample.

Equivalents

30 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

-55-

CLAIMS

The invention claimed is:

- A chimeric multivalent immunoglobulin (Ig) Superfamily protein analogue comprising one or more polypeptide chains forming a β-barrel domain containing complementarity-determining region-like (CDR-like) regions and framework region-like (FR-like) regions, said CDR-like regions defining a ligand binding site and said protein analogue having at least one additional ligand binding site segment spliced into the FR-like regions of said β-barrel domain.
 - 2. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein said polypeptide chains have an amino acid sequence wherein said sequence is substituted or modified in the amino acid sequence of at least one amino acid residue.
 - 3. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein a non-covalently associated two chain polypeptide forms a β -barrel domain.
- 20 4. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein a single chain polypeptide forms a β -barrel domain.
- A chimeric multivalent Ig Superfamily protein analogue of Claim 1 comprising a single chain polypeptide
 forming a β-barrel domain wherein said single chain polypeptide is comprised of two polypeptide chains connected by a polypeptide linker spanning the distance between the C-terminus of one chain to the N-terminus of the other chain.

- 6. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein the polypeptide chain is selected from the group consisting of: heavy chain (H), light chain (L), α chain (α) β chain (β), γ chain (γ), δ chain (δ), or ϵ chain (ϵ).
- 7. Biological material having a nucleotide sequence which encodes a chimeric multivalent Ig Superfamily protein analogue of Claim 1.
- A replicable recombinant DNA expression vector
 containing the nucleotide sequence of Claim 7.

5

- 9. A chimeric multivalent antibody analogue comprising one or more polypeptide chains forming a β -barrel domain containing complementarity determining regions (CDRs) and framework regions (FRs), said CDRs defining an antigen binding site, said antibody analogue having at least one additional antigen binding site segment spliced into FRs of said β -barrel domain.
- 10. A chimeric multivalent antibody analogue of Claim 9 wherein a non-covalently associated two chain
 20 polypeptide forms a β-barrel domain.
 - 11. A chimeric multivalent antibody analogue of Claim 9 wherein a single chain polypeptide forms a β -barrel domain.
- 12. A chimeric multivalent antibody analogue of Claim 9
 25 wherein said CDRs and FRs are comprised of heavy chain
 (H) polypeptide chains and light chain (L) polypeptide
 chains derived from variable regions (V) of
 immunoglobulin proteins.

-57-

13. A chimeric multivalent antibody analogue of Claim 12 wherein the CDRs spliced into FRs of the β -barrel domain to form an additional binding site segment such that a variable heavy chain (V_H) CDR is spliced into a V_H FR to form a V_H(V_H) polypeptide chain.

5

10

- 14. A chimeric multivalent antibody analogue of Claim 12 wherein the CDRs are spliced into FRs of the β -barrel domain to form an additional binding site segment such that a variable light chain (V_L) CDR is spliced into a V_L FR to form a V_L(V_L) polypeptide chain.
- 15. A chimeric multivalent antibody analogue of Claim 12 wherein the CDRs are spliced into the FRs of the β -barrel domain to form an additional binding site segment such that a V_H CDR is spliced into a V_L FR to form a $V_L(V_H)$ polypeptide chain.
- 16. A chimeric multivalent antibody analogue of Claim 12 wherein the CDRs are spliced into the FRs of the β -barrel domain to form an additional binding site segment such that a V_L CDR is spliced into a V_H FR to form a $V_H(V_L)$ polypeptide chain.
- 17. A chimeric multivalent antibody analogue of Claim 9 comprising a single chain polypeptide forming a β-barrel domain wherein said single chain polypeptide is comprised of two polypeptide chains connected by a polypeptide linker spanning the distance between the C-terminus of one chain to the N-terminus of the other chain.

- 18. A chimeric multivalent antibody analogue of Claim 17 wherein said two polypeptide chains connected by a linker further comprise two $V_H(V_H)$, $V_L(V_L)$, $V_H(V_L)$ or $V_L(V_H)$ polypeptide chains.
- 5 19. A chimeric multivalent antibody analogue of Claim 18 wherein to the N-terminal end of the polypeptide linker spanning the distance between the C-terminus of one polypeptide chain to the N-terminus of the other polypeptide chain is added a polypeptide residue bridge which connects the N-terminal end of the linker to the C-terminal end of a CDR sequence which has been added to the C-terminal end of a FR sequence.
 - 20. The polypeptide linker and bridge of Claim 19 comprising at least 19 amino acid residues.
- 15 21. A chimeric multivalent antibody analogue of Claim 9 wherein said CDRs and FRs are of mammalian origin.
 - 22. A chimeric multivalent antibody analogue of Claim 9 wherein said CDRs and FRs are of mouse myeloma origin.
- 23. Biological material having a DNA sequence which20 encodes the chimeric binding protein of Claim 9.
 - 24. A replicable recombinant DNA expression vector containing the DNA sequence of Claim 23.
- 25. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein, upon the binding of a ligand to one binding site, a conformational change is initiated in the β -barrel domain such that the affinity of the second binding site is modified.

-59-

- 26. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein an additional polypeptide effector molecule having a biological activity is linked to the N- or C-terminus of said β -barrel domain said
- 5 biological activity independent of the ligand binding activity of the chimeric multivalent protein analogue.
 - 27. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein one binding site is reactive with a diagnostic imaging agent.
- 10 28. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein one binding site is reactive with a radioisotope.
 - 29. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein one binding site is reactive with a cytotoxic substance.

- 30. A chimeric multivalent Ig Superfamily protein analogue of Claim 1 wherein one binding site is reactive with an effector molecule.
- 31. A chimeric multivalent Ig Superfamily protein analogue20 of Claim 1 wherein one binding site is reactive with a marker on a cytotoxic cell.
 - 32. A method for imaging specific tissue in a host comprising:
- a) administering to a host a chimeric multivalent Ig

 Superfamily protein analogue of Claim 1 having
 one binding site reactive with a targeted tissue
 specific antigen and a second binding site
 reactive with a diagnostic imaging agent under

25

conditions wherein said protein analogue binds to the targeted tissue; and

- b) administering the imaging agent to the host under conditions whereby said imaging agent binds to the chimeric multivalent Ig Superfamily protein analogue resulting in a detectable image of the targeted tissue.
- 33. A pharmaceutical composition for administration to a host for imaging specific tissue in a host comprising the chimeric multivalent Ig Superfamily protein analogue of Claim 32 in a pharmaceutically acceptable carrier.
 - 34. A method of irradiating specific tissue in a host comprising:
- a) administering to a host a chimeric multivalent Ig
 Superfamily protein analogue of Claim 1 having
 one binding site reactive with a targeted tissue
 specific antigen and a second binding site
 reactive with a radioisotope under conditions
 whereby said protein analogue binds to the
 targeted tissue; and
 - b) administering the radioisotope to the host under conditions whereby said radioisotope binds to the chimeric multivalent Ig Superfamily protein analogue wherein binding of said chimeric protein analogue to targeted tissue and binding of said radioisotope to said chimeric protein analogue results in irradiation of the targeted tissue.

-61-

35. A pharmaceutical composition for administration to a host for irradiating specific tissue in a host comprising the chimeric multivalent Ig Superfamily protein analogue of Claim 34 in a pharmaceutically acceptable carrier.

36. A method of delivering a cytotoxic substance to specific tissue in a host comprising:

5

- a) administering to a host a chimeric multivalent Ig Superfamily protein analogue of Claim 1 having one binding site reactive with a targeted tissue specific antigen and a second binding site reactive with a cytotoxic substance under conditions whereby said protein analogue binds to the targeted tissue; and
- 15 b) administering the cytotoxic substance to the host under conditions whereby said cytotoxic substance binds to the chimeric multivalent Ig Superfamily protein analogue, wherein binding of said chimeric protein analogue and binding of said cytotoxic substance to said chimeric protein analogue results in delivering the toxic substance to the targeted tissue.
- 37. A pharmaceutical composition for administration to a host to deliver a cytotoxic substance to specific
 25 tissue in a host comprising the chimeric multivalent Ig Superfamily protein analogue of Claim 36 in a pharmaceutically acceptable carrier.

- 38. A method of lysing target cells in a host having cytotoxic cells comprising administering to a host a chimeric multivalent Ig Superfamily protein analogue of Claim 1 having one binding site reactive with a surface receptor of a cell targeted to be lysed, and a second binding site reactive with a marker on a cytotoxic cell under conditions whereby said protein analogue binds to the targeted cell and said cytotoxic cell binds to the chimeric multivalent Ig Superfamily protein analogue, wherein binding of said chimeric protein analogue and binding of said cytotoxic cell results in lysis of the targeted cell.
- 39. A pharmaceutical composition for administration to a host a cytotoxic cell which lyses a target cell
 15 comprising the chimeric multivalent Ig Superfamily protein analogue of Claim 38 contained in a pharmaceutically acceptable carrier.
- A method of modifying the function of a cell surface receptor of specific tissue in a host comprising administering to a host a chimeric multivalent Ig 20 Superfamily protein analogue of Claim 1 having one binding site reactive with a targeted cell surface receptor and a second binding site reactive with an effector molecule under conditions whereby said protein analogue binds to the targeted tissue and said 25 effector molecule binds to the chimeric multivalent Ig Superfamily protein analogue, wherein binding of said chimeric protein analogue and binding of said effector molecule results in selective modification of the function of the targeted cell surface receptor. 30

-63-

41. A pharmaceutical composition for administration to a host an effector molecule which modifies the function of a cell surface receptor of a specific tissue comprising the chimeric multivalent Ig Superfamily protein analogue of Claim 40 contained in a pharmaceutically acceptable carrier.

5

- 42. The chimeric multivalent Ig Superfamily protein analogue of Claim 1 having one binding site reactive with a preselected ligand and a second binding site reactive with a substance labeled with a radioisotope or enzyme suitable for use as a quantifying agent in an in vitro diagnostic assay.
- 43. The chimeric multivalent Ig Superfamily protein analogue of Claim 1 having one binding site reactive with a preselected ligand and a second binding site having catalytic activity.
 - 44. The chimeric multivalent Ig Superfamily protein analogue of Claim 43 wherein both binding sites have catalytic activity.
- 20 45. The chimeric multivalent Ig Superfamily protein analogue of Claims 43 and 44 contained in a physiologically compatible carrier solution for administration to vertebrates.
- 46. The chimeric multivalent Ig Superfamily protein
 25 analogue of Claim 1 crosslinked at sites other than
 ligand binding sites to form a two dimensional array
 of chimeric multivalent protein analogues.


- 47. A method for producing a chimeric multivalent Ig
 Superfamily protein analogue comprising the steps of:
 - a) determining the splice points for CDR-like regions to form additional ligand binding site segments on the FR-like regions of a β -barrel domain whereby insertion of CDR-like region amino acid residues into the FR-like region residues maintains the folded structure required for binding activity with a preselected ligand;
- b) determining the amino acid sequence of the resulting construct having a first ligand binding site and a second ligand binding site;
 - c) deducing the DNA sequence encoding the amino acid sequence of b);
- d) synthesizing the DNA sequence;
 - e) inserting the DNA sequence into an appropriate expression vector and expressing the polypeptide in a suitable host system;
 - f) isolating and purifying the expressed polypeptide; and
 - g) refolding the purified polypeptide to its immunologically reactive conformation, thereby resulting in a chimeric multivalent Ig Superfamily protein analogue.
- 25 48. The method of Claim 47 wherein determining the splice points is accomplished computationally by use of a computer-generated three dimensional structure of the chimeric multivalent Ig Superfamily protein analogue.
- 49. The method of Claim 47 wherein determining the splice points is accomplished by primary sequence alignment.

-65-

- A method for effecting cell-cell interactions 50. comprising administering to a host a chimeric multivalent Iq Superfamily protein analogue of Claim 1 having one binding site reactive with a targeted cell surface receptor of a first cell and a second binding 5 site reactive with a targeted cell surface receptor of a second cell under conditions whereby said protein analogue binds to said first cell and second cell wherein binding of said first cell and second cell to the chimeric protein analogue results in interaction between the two cells.
- A molecular switch comprising a chimeric multivalent 51. Ig Superfamily protein analogue of Claim 1 having one binding site initiating a conformational change in said chimeric protein analogue when said binding site 15 is bound to ligand, whereby the conformational change causes the chimeric protein analogue to act as a molecular switch.

- A chimeric multivalent Ig Superfamily protein analogue 52. according to Claim 1 for use in therapy or diagnosis. 20
- An analogue according to Claim 52 for use in (a) 53. imaging specific tissue in a host; or (b) irradiating specific tissue in a host; or (c) delivering a cytotoxic substance to specific tissue in a host; or (d) lysing target cells in a host; or (e) modifying 25 the function of a cell surface receptor of specific tissue in a host; or (f) effecting cell-cell interactions in a host.

- 54. Use of a chimeric multivalent Ig Superfamily protein analogue according to Claim 1 for the manufacture of a diagnostic agent for imaging specific tissue in a host.
- 5 55. Use of a chimeric multivalent Ig Superfamily protein analogue according to Claim 1 for the manufacture of a medicament for (a) irradiating specific tissue in a host; or (b) delivering a cytotoxic substance to specific tissue in a host; or (c) lysing target cells in a host; or (d) modifying the function of a cell surface receptor of specific tissue in a host; or (e) effecting cell-cell interactions in a host.



SUBSTITUTE SHEET





FIGURE 2A



SUBSTITUTE SHEET

000000000	†			
000010000	: :			
4444444	1 1			
33333333	 81			
HZHKHHZH	. H-	- !		
4444444	i			
	+ 4	aaaaaaxt	:	!
SKKKSKKA		1		i
ZZZOHZZZZ	•	REMERKE		ı
M I W D Z D O H I		PPPPPH	ISI	J
OI I I O I I H W	!	******	H -	_
Z I I O I I A O		MXXXXMX		
о ттитин	, ,	* ************************************	·	1
αιιαιια	1 10 CD RD	ואצצטטא	Ë	i
Z I I I I I I I I I	្រដ្ឋប	, 0 44444	C	,
コエ・マン・・・	į Ei	០ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១ ១	-	_
дннідініі	30	8 11 11 18 18 18 18 18 18 18 18 18 18 18	30	
a x U a a a a a i fi		। मधामधाम्य		
0000000000		ниндный	17.	
α α α α α α α α α		(四本草草木草草	i	
O A A A O A A A O	i	0000000	į	
K R R S S S S S S S S S S S S S S S S S		ខ្លួលខ្លួ ង ម	į	
0.00000000	1	HOAAAA	i	
α \exists \exists α \exists α \exists	2	AHKAPK		
жинни ж анн	80	0000000	į	
8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8		 Q 		
>>>><	2	анхаана	20	
RHKKOKOKK		ROKKKKO		
日日日日日日日日日日日		4424224	į	
11811111		លលលលកល ក	•	
000000000	II.	0000640		
A S W H H H H H A M A M		0 0 0 0 0 0 0	નું	
ស ជា 🚓 ជា ជា ជា ជា ជា ជា		на фини	BL1	
>404>4>40		OKKOOKK		
BONDAPHOO		484444		-
ายผลเสสาน เ			P	i
и и и н и н и и и	0 4	្	0 1	
MAHAHAMA		0 A A O O A A	-	
РЕВРЕМЕН	80	000000	1	
ល្យ ស្រុក្រលួល្ប ្		ფფფთ ₽ ფფ	ੂਜ਼ੀ	
			80	
		ן ממטניאאל	1	
- REELEREE		444444	1	
>00>>>00	1 1	ROOKKOO		
нннн>ннню		222222	:	
еремереем (i	почины	ed :	
	· · · ·			-
д м м м о м о н ы і	ם מח	1 1 2 2 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	osition Regions Kabat	i
。 O D 4 D 4 展 B (O H 6 D 7 B 1 B 1 B 1 B 1 B 1 B 1 B 1 B 1 B 1 B	tion	א ש ש יי ו ש ש	1 t t t d	
MCPC 1 R.1 HYH HYH	4 9 3	10Pc D) R10 4-4- HYRe	81 99	í
MC 44 H H H H Y	ğă	H Y	90 90	
	-	##### ##	+4	

FIGURE 2B


•			
		наниина	80
		លលក្ខពុធ	w :
		OKNKKNK	1
		លលល≮លលល :	- F
		HZKZAHH	į
		дааааа	i
		RKDKKAK	
		***************************************	84
	. 1	нанны 4	ö
GGDaDaDGa		навнивы	i
*****	HI H	H P M M H H M	2-1
$\mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} \mathbf{o} $	F	RKKKKK	
ω	i i	0000000	
0000000×		KKKKKK	3
ω ω ω ω ω ω ω ω ω		1 444444	BL
00000000 0 4	084	84 4 8 8 8 8 9	
F	2	4000000	
14.14 14 14 14 14 14 14 14 14 14 14		OZZHOHZ	
我我我我我我我我			ູ່ຕຸ
O a a K O > a a O		ROWENER	OSS: CDR
приприпри	eg i	наннын	8 5
>>>>>>	B.L.	RENHEROR	
$ \verb"o" o o o o o o o o o o o o o o o o o $	<u> </u>	סטאטטא	· · · · · ·
$\alpha \cup \alpha \alpha \alpha \alpha \alpha \alpha \triangleleft \alpha$		RDKSZGS	
国内耳丸平丸TOP		ZIOCHOI	~
ませせませるせせ	083	σιιρ ιτί	TL2
HEKKZKOZJ		KILIKIT	
ល H ល ល ល ល ល ល ប	912	*****	1
444444	<u> </u>	WENERE SE	
よ耳よ口 ¥ 耳えより		мннннь і	ı
先在 第五五五五五五		よぼりほよ対す	22
нььннннн	21	ৰ ৩ ৩ ৩ ৰ ৩ ৩	
444848444	i Öi	т қн 4 н ң н	1
はせせるひませむせ	i i	KEEEEEE	1
KOKKKKKKK	<u>i i</u>		
84 14 14 14 14 14 14 14 14 14 14 14 14 14	i	1 444444	1
ው ል ዙ ዉ ል ል ል ል ፋ ፋ	101	K G G G G G K i	
OKOHOHEKK	1 2	KKOKK#X ¦	BL2 2
o o o o o o o o o	, m	២២២២២២ ។	m c
ч от ат а а т т	i pr	ррррррр	FR
K K K K K K K H >	<u> </u>	<u> </u>	64!
		40 00 00 00 00 00	44 44 44
жиге - бор	こははせ	1123.00 10509934	sitions Regions Kabat
CPc60 D1. R19. 753 753 HYHEL YHEL1	trion don aba	00174#0 0498194	osition Region Kaba
CPC R3 R4-4 KHIX	1 4 6 X	MCPC R1 A - 4 HYH	B X
MCP R 4- HYH	្តែស្តី	ž Tr	Po
보보보보보보보보	-	****	

FIGURE 2B

•		ທ ∢ ຸ
		တ တ လ လ လ လ
		нананны
		464464
		ннимнин
		H H H H H H H I O 44
		0 0 0 0 0 0 0 1 H 4 F
& >		୍ ଶ୍ରସ୍ଥ୍ୟର । ସ
KKKK KKHH	! !	
нннаннна		******
моммами	t !	ו ואטאאמאל
4444444		9994999
KKKKKKKKO		0432464
04444444	4	וםטאאאא
000000000	84 84	וואאטאאצ
40000000	ĤŢ.	нарынно
000000000	į	מולאווו ול מ
44444444		ριηιιιι (Ε΄)
	i i	>
84448HXPH		
наяханая		1101111
किस्ट्रा देशसम्ह	ام	
SOUTHEROUD	(4.5)	
ZOZUHAJINI ZOZUHAVZH	SEB BEB	ОККЖАВК
24020202 240202020		Z
ZHOOOOOA		ZEGRONZ O
00000004		TI PERPAPA
000000000	'	0000000
20000000	<u> </u>	(M I
	HS	B
*************************************	<u> </u>	нкрань
404404044	i i	4444004
инарернаг	 i	HENHENH
00000000		дааааа
APHACHHO	4.0	4 H W W D W H P H W H W
	8 8 8	M M M M M M M M M M M M M M M M M M M
PP4X4XPP9		444444
N N Z H K N N N O		4008200
w z w z w w z w w	 i	ZZKWZZZ
идидиидии ннининни	i	zzazzaa:
		akaaaaa wi
PHRHXHRAH	S	1424424 0 1
		-
Haaabaha		t t
マスゴスゴスススゴ	 i	
		VH PD 03: PD 03: PD 03: CD 03: CD 03: CVH 015: TH 01: CH 10: CH 1
SO MADEME		WCPc603: D1.3: R19.9: U539: 4-4-20: HYHell0: Ositions Regions
200 100 100 140 181 181 181 181 181 181 181 181 181 18	Tti Xio Xio Cab	Poble Hold Residence of the Pople Residence o
MCP R 4- HYH	: 88 : 48 : 48	MCP R 4- HYH 081
	, 0 tt	ρ.,
чччччччч		


IGURE 2B





SUBSTITUTE SHEET





SUBSTITUTE SHEET





FIGURE 4A



SUBSTITUTE SHEET






FIGURE 4B



SUPATIONALOWERT





SUBSTITUTE SHEET

FIGURE 4C




IGURE 4D




SUBSTITUTE SHEET






SUBSTITUTE SHEET

TROKE 4E








SUBSTITUTE SHEET








SUBSTITUTE SHEET






Aligned Pos. #			H102				H114											
•						ŧ								ŧ				
н3	YY	C	A	R	N	Y	Y	G	S	T	W	Y	F	D	<u>v</u>	W	G	A
								*										
		*	*	*	*										*	*	*	*
H BL2		W	v	R	Q	P	₽	G	K		R		L	E	W	I	A	A
Conserved Rea	в.	*		*	*									*	*	*		
					Ť										•			
Aligned Pos.	#				H.	39									H	47		

FIGURE 6

SUBSTITUTE SHEET




Splice Points

Aligned Pos.	#						L!	99					L	104	1		
L3	Y	Y	С	Q	N	D	H	S		Y	P	L	Ţ	F	G	A	G
	#	*	•	*	*	*	-				-	-		*	*	*	
L BL2 Conserved Res		A	W	Y	Q	Q *	K	P	G	Q	P	P	K	L	L	I	
Aligned Pos.						† L	45							† L!	53		


FIGURE 7

SUBSTITUTE SHEET



Aligned Pos. #	H31						H:	H33			
H1 ·	F	T	F	s				M	B	W	
	*	*	*	*	*	*	*	*	*	*	
H BL4	R	A		_	T	A	I	Y	Y	C	
Conserved Res.			*	*				*		*	
Aligned Pos. #				H!	92			H!	96		

FIGURE 8



Splice Points





FIGURE 9







FIGURE 10






FIGURE 11

L2'

22/35

FIGURE 12

McPC603(McPC603): $V_H(V_H)-V_L(V_L)$ χ -Protein

N-terminus: H FR1 EVKLVESGGGLVQPGGSLRLSCATSGFTFS HI DFYME H FR2A WVRQ YYGSTWYFD H3' H FR2B ASRNKGNKYTTEYSASVKG H2 H FR3A RFIVSRDTSQSILYLQMNALRAED HI' DFY H FR3B YYCAR **H3** NYYGSTWYFDV H FR4 WGAG H2' SRNKGNKYTTEYSASVKGRAIV bridge AAAAA linker SGGGGSGGGGGGSL FRI DIVMCQSPSSLSVSAGERVTMSC - L1 KSSQSLLNSGNQKNFLA L FR2A WYOO **HSYPLT** L3' L FR2B LLIY L2 GASTRES L FR3A GVPDRFTGSGSGTDFTLTISSVQAEDL Ll' LLNSGNOKNF L FR3B YYC QNDHSYPLT L3 L FR4 F G

YGASTR ESGVPDC

			·
-	aaaaaax	Xaaaaaaaaaa	00
	KKKKKK	*********	KK !
	>>>>>	н>>>>>>	Si re
	3333333	ZZZZZZZZZZZZZZZZZZZ	33 Hi
	MEZNZHN	ZHKHHUNZ N Z >	ZI
	E>>EHE	PHZZZZZZHH	ΣH
	KEEEOOK	りりゃぬぬええぬえるえ	ਮ ਮ ਜ਼ੂ
ने व	ดิหหหหหล	み れれよよなよよれの	64 M
#	ัดยหมดดห	ανονασακκονο	ا یم
	NEENNNEE	EFFEX00000E0	EHO !
	начачаца	로	66.
	6049940	លលម្លងម្នងម្	HE 끝
	ロスよよなよ	れたよりよみはよれよは	×× `
	0000000	00000000000	00
	សលលលលម	Θ	လ လ
	HDAAAAD	>> A A A H A A A A A	νA
_	4545	WHXXHA4>>XX	X E4
	0000000	00000000000	υυ
	SHOOOOH	ϵ	တတ္က ျ
	บหมายหา	ㅋㅋㅋㅋㅋㅋㅋㅋ	хнойі
	ROXXXXX	500222222200	~~ "
	ちちかなれなり	4422244242	>> i
_	មេខមេលលល ស	\mathbf{R} \mathbf{N}	SS
	0000440	QQ4440000000	44
	0 0 0 0 0 0 0 0	ოოიიიიიიიი	100 31
	PUKUUUU	66666666666	교 교
	OKKOOXX	KAKKOOOKKK	i X M
_	>>>>>	>>>>>>>>>>	i > >
	нананан	аныныныны	नन
	0000000	្ត្រ	
	0 P K 0 0 K P	4444666644	44
	0000000		100
	SOSTOS	ι ωωωμωωωωωωω	2 2 S
	E TE		i
	> x x y D a a	QXQQQ>HB>QQ	100
	- нананан	даааааааааа	144 1
	* OOX * OO	aaaaaxxx xaa	100 1
	>>>>>>	>>>>>>>>	> >
	E C C E E E E E E	попопининия	M O H
 0 5		88666666666	0 4 E 0 t
in the	10000000000000000000000000000000000000	HHHHHHHH > >	2610 LOOP4 ition gions Kabat
þJ	こりょりゅばる	ннннн	
X-splice	MCPCPC R HYHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHHH	0 0 0 0 0 0 0 0 0 0 0 0	oα
×	.,	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	Д

FIGURE 13A

والمراوي والمتماعين والمراوي والمراوي والمناوي والمناوي

1	•		
	Į.		
i	s «	Κ ΩΝΩΝΩΚΩ Ω	ω ω •••••
		+	<u> </u>
ار	>>>>>	4 4444444	88
E	444444	14H 444444	>1
H2'(T	5445444 5445545	TORRORYY BY	v ⊨
-	444444	на нанинан	400
	0000000	00000000 00	7 4 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
	4000000	00400400 00	H O H
	0000000	00000000 00	00
	******		33
	>>0>>>	טא אאלאאלאא	××
		40000044000	۵۵
	CHEZZEG	ጀመሪያ ከሚከተር ነ ታ	ጀμ
	וםטאאאו	エロマニカネザススニー	« >
	ואאטאאצ	אראוט צם מצוו	≯ ≻
	1114404	ממאו איא מסוו	X X O
6	מולאווו	1110KW>1HOH	द्रात ह
<u>:</u>	י יוווהוט	וטאוומהאווו	. 요 1 4 표
H3'	וווומוא	ן ז אל או טו או וויי ו	- 1 1 월
	1 1 10 1 1 1 1		
	1 10 1 1 1		
	110111		
	1131111	KK	ν _Ε
	_ X X Y X X X X X X X X X X X X X X X X		ς ×
		OKKKKKKHKKY	
	AAAAHAA	HAAKAAAHAAA	0 1
	0000000	00000000000	
	**************************************	 44666666	n 1
	************************************	***************************************	ו אא
	H K P H H P H I	444444444	>>
1	FRAKABUSA	<i>~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~</i>	AA
	1 6606706	80000000000000000000000000000000000000	တ တ
_			00
	២០២២២២២	момымымымы	4.U
	4 H W W > W H	1	FR 90 S S I
		HOHHHMMMMHM	}
		> + + + + + + + + + + + + + + + + + + +	
	ANNXXNN	t	·
	ZZKOZZZ	1	1
		1 42444222244	I I
		0x00000000	
	**************************************	XXTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT	l t
		1	1
		· · · · · · · · · · · · · · · · · · ·	+
0			10: P4: ons bat
Č	7 2 3 3 3 4 4 6 4 6		2610: GLOOP4: sitions Regions Kabat
G	7 C C C C C C C C C C C C C C C C C C C	і нананні і нана	40 th 22 x
U.	MCP R A HYH	000000000000000000000000000000000000000	U
\sim		συσυσυσυσο	<mark></mark>

FIGURE 13A

•	нанимна	аоннинихне	H H O.
	νννΣννχ	Z000Z0Z0Z0Z	ທທີ່
	OKNKKNK	XXNNNQXXXNH	ss H
	νανκανα	α	ທທ ີ່
	HZKZDHH	HXXXHHXAXH	××
	999999		ا مما
•	RX>KKAK	AKANANA KAKA	> <
	HHRRHRR		HH 4
,	наннын	ннаанриннан	L L
	оннным	омененненен	HE !
	нучный	нымамымынь	440
ŀ	RKKKKK	REXXXXAXXXXX	XX
1		0000000000	00
	000000	•	1
ı	XXXXXXX		F
- 1	> 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		×z
- 1	SAKSSAS	<u>.</u>	αμ
	ANDUDEN		·
	ひとれ 子 ひ 耳 と	ZZZZOWHKAZK	22
ار	まままままま。		ם אאן
S	KKKKOOM		S KG
'n	наннын		HH0
H	ちらままれれよ	XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX	> =
	שטאטטטא	> O O O O O O O O O O O O O O O O O O O	00 🛱
Ì	KOKOKOO	αυχαχανχαχα	ωz
	וטאטוא	NAZZAZOEOO	ן א טאן
	0111411		[나]
- 1	XIIIXII	11111111111	[
- 1	200026	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	ሰ ሰ
ı	ኢንአቷሊካ Ω	NZZDOKZKUZU	ا حدا
	«ннннн»	ннннюннннн	інн
	よ 至 り ほ よ は マ	えたよぼほびなななん ス	ં×≭દ્ધું _
	4000400	७७७७७४७४७७७	ပတ္မွ
	ичнньия	でまんんままますで就	нн 🗀
	KEEEEEE	OSSSSSSSSS	×≈
<u></u> i	២២២២២២២	២២២២២២២២២២២	QΜ
		дыынынынын	i a a
£	αυυυυυα	KONOOKOOKOO	ပလ
E),	XXOXXEZ	ZXXKOXXXXOO	ן אַס קי
EH.	០០០០២០០		്രം ജ്
	444444		E E
	PPRANKE	FRORKARANEKA	1 0 20 1
		· · · · · · · · · · · · · · · · · · ·	+ 1
61	# m m o o o o o		8 E E E
ŭ	2016 1016 1016 1016 1016 1016 1016 1016	HHHHHHHH+>>	2610; GLOOP4; seltions Regions Kabat
ŭ	CPC6 D1 R19 J5 J5-4-4-HYHe		26. GLOOJ Sitio Regie Kal
X-splice:	MCPC6 D1 R19 J2 4-4- HYH6		OS J
×	بد بد	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	ŭ

IGURE 137

13B
RE
GG
댐

H

-	DODNONDON	ADONNOX	AA
	нанананы	2999999	G G G G G G G G G G G G G G G G G G G
	000000000	0020000	ပေပ 🛱 📔
	NNNNNNNN	ναναναΗ	တတ
-	<u> </u>	4000000	ပပ
	ναναναναν	ννννννν	ດດ ີ້ບຸ
	იიიიიიი	0000000	P. S. S. R.
	μονονονοι	F1 00 00 00 00 00	N N O]
	म म म म म म म म म	存在存货货货	E4 E4
 	KKKKKKKK	RRKKKKK	**
1	∪ 000<0>000	DOAKKOK	00
-	ририририи	припини	2 P B I
- 1	>>>>>>	>>H>>>>	
	000000000	0000000	00 <u>i</u>
<u>S</u> .	SOSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSSS	HOOOOOA	SO
•	MAHAMAHOA	VEBKKEK	i fe fe
Z,	スセロススという	KKTTTTK	KK 0.4
- 1	HERKKKKOKH	ZZZZXXZ	1 Z Z G H
ſ	N F1 N N N N N N D		V S V S TIL2
	ZPPHCHHP	ADARHER	+ 1
- 1	**************************************	OKKKRYO	××
_1	ままままままま	DKKKKKK	~ ~
	нььннннн	ннннннн	H H H
	นหษายนะ	प्रमाद्रद्रमम	1
	なななななななな	ה אשר אים ו	145
	XOXXXXXXX	XXXXXX	† * * *
	₽₽>₽₽₽₽₽₽	1 2 2 2 2 2 2 E	וְאַהָּה מִי
£)	4240000044	THE WORLD	FR 50.1
13,	QXGEQEBXX	IOOOKHUE	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
ä	, 000000x00		1 1
	TOTOTOTOTO	 AAAAAAA	A A
	XXXXXXXX	<u> </u>	†**
••		i	
9	VL 603 11.3 15.39 15.39 17.20	P C C C C C C C C C C C C C C C C C C C	2610 GLOOP4 Sition egions Kabat
7	CPC603 CPC603 D1.3 R19.9 7.539 VA = L-20 XHEL10 REI10	LAAAAAA	2610 LOOP4 ition gions Kabat
S	VL MCPC603 D1.3 R19.9 J539 4-4-20 HYHELS HYHELS REI	KAPPA KAPPA KAPPA KAPPA KAPPA KAPPA	2610 GLOOP4 Position Regions Kabat
X-splice	E	KAPPA KAPPA KAPPA KAPPA KAPPA KAPPA KAPPA	ı og
• •			

FIGURE 13B

	A	1
	. < 0	
× >	RKKKKKC	~ ~
HHKK KKHH	XXXXXX	××
нннаннна	чнннчн>	IHH !
5, инининия	ымымымы	មក
🔑 មកម្មមាមមុខ	нананана	HH
y KKKKKKKO	*****	XX '
оныныныны 💾	нененне	
00000000	0000000	50 0 E
4004000A	ΑυνΑΑύ υ	6644.
იიიიიი	0000000	00
ા માં માં માં માં માં માં	4444444	44
	 6666665	HH
ם א א א א א א בו	назчнка	1 1
O DARAHAZARA	TULADAT	1 2 2
DYKYNAHWNC ()	ZHAADAK	
L NOFXERNOU!	OKKEKO	## 4
**NHHUZOZ	KZZZZAK	EJ CJ
OHUBWBW N	E U E N N N H	600
	1113111	1 1 2
22000000A	aaaaaa	00
00000000	40000A	0 84
00000000	000000	00 7
***************************************	BBKKKBK	13 KH
 	**********	H KK
	нання	нь і
HADAGAGAA	HUKKKKK	00
ны на на на на	HHADKHE	1 200
00000000	000000	00
តាមាតាមាតាមាតា	ммоммим	មេខ 🕴
APHAMHHAN!	>444408	44 30
оспыныноы ј	онинино	E E E
והרלבלברים	>>>==	>> 1
លលកម្មលលល ា	ναμνναυ	KK
OZOZOOZOO	BONDENO	0 0
і нанананы і	ннннны	нн 1
HXHHXHWHA	нинненн	×× 0 !
	444444	1100 8
	HHHOOOAI	H H
	VHKKHHH	ELE I
	""H""""	04581
11.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0	HHHH	2610 00P4 tion ions abat
X-splice VL MCPc603 D1.3 R19.9 4-4-20 HYHELIS HYHELIS REI	PPA PPA PPA PPA PPA LAMB	2610 GLOOP4 Position Regions Kabat
A H H	KAPPA KAPPA KAPPA KAPPA KAPPA KAPPA	ဂ လို ရှင်
×	XXXXXX	Ā ⁻

FIGURE 13B

MCP(MCP): EVKLVESGGLLVQPGGISLRLSCANTSGFTFSDFYMEWVRQLYGBTWYFDR19(D1.3): QVQLKESGAELVAASSSVKMSCKAASGYTFTSYGVNWVKQRDYRLD 2610(D1.3): EVQLQQSGPELVKPGASVRMSCKSSGYIFTDFYMNWVRQRDYRLD 610(GLOOP4): EVQLQQSGPELVKPGASVRMSCKSSGYIFTDFYMNWVRQTHYYFD Regions: Regions: Kabat:: Kabat::	MCP (MCP): W I A A S R N K G N K Y T T E Y S A S V K G R F I V S R D T S Q S I L Y L Q M N R19 (D1.3): W I G Y I N P G K G Y L S Y N E K F K G K T T L T V D K S S T A Y M E L R 2610 (D1.3): Y I G Y I S P Y S G V T G Y N Q K F K G K A T L T V D K S S T A Y M E L R 2610 (GLOOP4): Y I G Y I S P Y S G V T G Y N Q K F K G K A T L T V D K S S S T A Y M E L R Regions: IS2	MCP(MCP): A L R A E D D F Y Y Y C A R N Y Y G S T W Y F D V W G A G B R N K G N X Y T T E Y B A B V K G R I S I Z E D G Y G Y F C A R S F Y G G S D L A V Y Y F D S W G Q G I W G D G N T D Y N S A L K S R L S I Z E D G Y G Y Y Y Y C A G S S G N K W A M D Y W G H G I W G D G N T D Y N S A L K S R L S I Z I Z E D N Y Y Y Y C A G S S G N K W A M D Y W G H G I Y P G N G N T K Y N E N F K G K A T L Regions: Regions: Regions: Rabat: H FRJA H FRJB H FRJB H FRA H	MCP(MCP): A A A A G G G G G G G G G G G G G G G
R1 2610 2610 (M R1 2610 2610(M R1 2610 (R1 261 2610 (

FIGURE 14A

· .3

1,3740




FIGURE 14B

SUBSTITUTE SHEET

```
AGICTCTGGTCCATTCTAATGGTAACACTTACCTGAACTGGTACCTGCAAAAGGCTGGTCCGAACCTTCTGATCTACAAAGTCTTAAAAGTCTTCTGGTGTCCGGAACTCTT

Q S L V H S N G N T Y L N Y L Q K A G Q S P K L L I Y K V S N R F S G V P D R

LWNI:6 Avail:1

MACCGS:1

Hindili:1

Hindili:1

BSpM1)+
                                                                                                                                                                                                                                                                                      130 140 150 160 170 180 190 200 210 220 240
AGECTCATGGTAAGTCTCTAGACTACATGGGTACATCTCTGGGGTACAACCTACAAGTTAAGGGAAGCGACCCTTACTGTCGACAAATCTTCCCAACTG
Q S H G K S L D Y I G Y I S P Y S G V T G Y N Q K F K G K A T L T V D K S S S T tX1:8 Xbaldam:1 R$al:b PfINIdcm:7 BstEll:1 Drai:b Hincli:b
                                                                                                                                                                                                         CCATGGAAGTTCAACTGCAACAGTCTGAATTGGTTAAACCTGGCGCCTCTGTGCGCAATTCCTGCGGTACATTTTTCACCGACTTCTACATGAATTGGGTTCGCC
                                                    Rsal:b
                                                       Fsp1:b
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          FIGURE
                                                      Bsall:2
                                                                    Ehel:b
Narl:2
                                                                                                                                                                                                                                                                                                                                                                                                                                                                             BspM1)+
EcoN1:5
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        #3048- [VL-26-10] -up-
                                                                                                                                                                                                                                                                                                                                                                          BsaH1:2
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         Kpn1:5
Rsa1:b
                                                                                                                                                                                                   ====#3617===
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         CCAGGTCGAGATTAAACGTTAACTGCAG
T K L E I K R *
Avai:1 Hincilib
                                                        Avail:1
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              Pst1:5
                                          M E V Q
Ncol:1
Styl:1
                                                                                                                                                                                                                                                                                                                                                                            BspEldam:1
pH1560.seq
10
```

FIGURE 15B

120 GTTCGCC V R Bs	240 TCAACTG S T	360 360 361AATA 6 N	480 STCTCTGC S L	600 SAAGCTIC K L Hindill	720 TATCTACT 1 Y	078
110 ATGAATTGG M N U	230 CAAATCTTCC K S S 1:b	0 350 360 1GT 350 360 1GT GGGGGGGGTGGTATATA G V G D G N	470 GACTCCCGTGTCI T P L S Drdi:7 Nsp811:b	590 TCAGTCTCC	710 AGACCTGGG : D L G	830
100 CGACTTCTA(D F Y	220 TACTGTCGAC/ T V D Hincil:	340 TCATGGTGT H G V	460 AATGACCCA H T 9	580 AAAGGCTGG K A G	700 CGAGGCG6/ / E A E	820
90 CATTTCAC(1 F T	210 IGGCGACCCT	330 XTATTGGGG Y W G	450 CCGACGTCGT S D V V I Aatli:5 I:1 BSAH1:2	570 GGTACCTGCA W Y L Q Acc65:1 BspM1)* EcoN1:5 Rsa1:b	680 690 CCTGAAGATCTCTGTTCG L K 1 S R V Bg111:1 Bsty1 #3048- [VL-26-10]-up-	810 GGATCC BamH1:1
80 CCT CT GGGTA S S G Y RSB	40 150 160 170 180 190 200 210 220 230 240 TAGACTACCGGGACCTTACTCTCGGGGTTACCGGCTACACAGTTAAAGGTAAGGGACCCTTACTGTCGACAAATCTTCCTCAACTG L D Y I G Y I S P Y S G V I G Y N Q K F K G K A T L I V D K S S S I aldam:1 Rsalib PflHidcm:7 BstEll:1 Dralib Dralib Sali:1	==#3617== ==================================	440 450 460 AGCTCGAGTGGTAATGACCG S S G S D V V W T Aval:1 BstYl Aatli:5 Xhol:1 BamHi:1	S60 ACCTGAACT Y L N	680 ICCCTGAAGATC T L K 1 Bg[1] BstY] #3048-[VL-	790 800 810 GITAACIGCAGAAGCTTATCGGATCC R H HINGII:b HINGII:18amH:1 Hpai:b 8stYI
70 rectecaaate s c K :	190 AACCAGAAGT N Q K D	310 5GTAACAATI G N K 15281- [VH-	430 36TTCCAGCT 6 S S Av Av	550 GGTAACACTT	670 SGTACTCACTTC/ G T D F RSel:b	790 R * Hincilib H Hpalib Psti:5
60 retececater V R H Fspl:b	180 TACCGCTACA T G Y EII:1	300 :GGGCTCTCTC (G S S) 4 anti#3407-	420 TCATCTAGG(S S S dam:b 1	540 CATTCTAAT H S N 11:1 3stx1:8	660 16611C1GGT G S G Rs	780 GATTAAACGT 1 K R 1 H HI 1 HP
50 GGCGCCTCT G A S BsaHI:2 Ehel:b	170 ACTCTGGGGTT Y S G V Idcm:7 BstE	290 ACTATIGCGCG Y Y C A al:b	410 420 AGCTCCCGGATCTTCATCTV S S G S S S act:Ect36 BaseBldam:b BspEldam:b BspEldam:anti#3107-BASSlink-[#	530 GTCTCTGGT(S L V WMI:6 Ava HIdcm:7	650 CTCTGGTTC S G S	770 780 GCACCAACCCGAGATTAAA G T K L E I K AVBI:1
40 GGTTAAACCT V K P	160 TTCCCCATAC S P Y PflHIC	#3617== 280 TTATGGGTA(TY G Y F G Y RSa	400 ATCGAGCTC S S S mm:2Sacl:E SRV:b	520 SCTCTCCCA \$ \$ 9 A I Pfl	640 CGGATCGTTI	760 6 G G G T
30 IGGTCCTGAATT G P E L Avall:1	150 160 NTCGGGTACATTCC I G Y I S Rselib	270 280 270 280 7.CIGAGGACGGITATGGGTA S E D G Y G Y RSB	AGTCTAGACTATCGATATCGA AGTCTAGACTATCGATATCGA K S R L S I S Claidem: Sa Xbaldem: 1 ECGRV:b	510 117C77GCC0 1 \$ C 1	630 fcr66fercc s G V	750 :GCCGACTTTG
20 SACAGTCTGG Q S G AV	140 CCCCTAGACTACA1 S L D Y I Xbaldem:1	260 TTCTTTGACCT	TGAGTCTA	00 AGGCTTCTA 0 A S 1	620 AACCGCTTCT N R F	740 750 760 CATGINGTGGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTG
CAACTG(AAGTCTC K S Xb	2 ictgcgtt L R	### ##################################	490 500 510 520 530 540 550 560 570 580 590 600 CGTTTCTCGGGGGGGGGGGGGGGGGGGGGGGGGGGGG	610 620 630 640 650 660 670 680 690 700 710 720 TGATCTACAAAAGTCTCTAACCGCTTCTCTGGTTCTGGTACTGACTTCACCCTGAAGATCTCTGTGTCGAAGACCTGGGTATCTACT L I Y K V S N R F S G V P D R F S G S G T D F T L K I S R V E A E D L G I Y RS81:b BG11:1 #3048-UL-26-10]-up-	SO CAGACTACTO
pH1539seq 10 CCATGGAAGTII M E V NCOI:1 Styl:1	130 AGTCTCATGGT Q S H G tXI:8	250 CTTACATGGAG A Y M E	EEEEE#361 37 CCGATTACA T D Y	45 CGGTTTCTC P V S	61 IGATCTACA L I Y	7. 101661616 F C S

WO 93/23537 PCT/US93/04338

33/35

OXIDIZED x-1

REDUCED x-1




FIG. 16A

SUBSTITUTE SHEET

PCT/US93/04338 WO 93/23537

34/35

OXIDIZED x-2 MIXTURE

OXIDIZED 26-10 sFv




FIG. 16B




FIG. 17A




FIG. 17B




FIG. 17C

PCT/US 93/04338

		• • • •		International Application No	PC	1/05 93/0433
L CLASSII	FICATION OF SUBJ	ECT MATTER (If several classifi	ication sym	sbols apply, indicate all) ⁶		
According	to International Paten	Classification (IPC) or to both Na	tional Clas	ssification and IPC		
Int.C1	. 5 C12N15/1	2; C12N15/13	3;	C12N15/62;		2P21/08
	G01N33/4	8; GO1N33/5	34;	A61K37/02;	A6 :	1K39/00
II PIFI DS	SEARCHED					
		Minimum	Document	tation Searched		
Classificat	ion System		а	lassification Symbols		
Int.Cl.	. 5	C07K ; C12N	;	A61K		
		Documentation Search to the Extent that such Doc	ed other th	an Minimum Documentation e Included in the Fields Searched ⁸		
			-		. 	
III. DOCUI	MENTS CONSIDERE	D TO BE RELEVANT ⁹				
Category a	Citation of D	ocument, ¹¹ with Indication, where	appropriate	s, of the relevant passages 12		Relevant to Claim No.
X		ENGINEERING	- 110'	4ND 00		1,2,6-9,
-	vol. 5, no. 3, April 1992, ENGLAND GB				12-16,	
	pages 229 - 234 SIMON T;RAJEWSKY K; 'A functional antibody					21-25,
	21WOW I	;KAJEWSKY K; 'A Tur with an insertion i	1CT10N	al antibody framework		43,44, 47-49
		With an insertion i 3 loop of the VH do				47-43
1	region .	tions for antibody	engin:	eering !		
·	implications for antibody engineering. see the whole document			1	3-5,10,	
					11, 17-20,	
	:'					26-42,46
			-	•		
·				-/		
. "						
ł						
		•				
		-				
° Special	categories of cited do	coments: 10		T' later document published after th	e Internat	tional filing date
"A" doc	ument defining the ger	eral state of the art which is not		or priority date and not in conflicted to understand the principle	ct with the	e application but
COL	sidered to be of partic	ilar relevance		Invention		
	lier document but publi ng date	ished on or after the international	•	"X" document of particular relevance cannot be considered novel or ca		
"L" doca	ument which may thro	doubts on priority claim(s) or		involve an inventive step		
	which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relevance; the claimed inventive step when the considered to involve an inventive step when the considered to involve step when the constant step when the					
		oral disclosure, use, exhibition or		document is combined with one of ments, such combination being o	or more ot	her such docu-
	er means ument published prior	to the international filing date but		in the art.		
	r than the priority date		•	'A" document member of the same p	atent fam	ily
V. CERTIF	FICATION					
Date of the	Actual Completion of t	he International Search		Date of Mailing of this Internation		th Report
	10 SEPTEME	BER 1993	1	28-09-	1993	
	TA API I FUIT					
nternational	Searching Authority			Signature of Authorized Officer		
	EUROPEA	IN PATENT OFFICE		NAUCHE S.A.		
		·				

, 2

Y BIOCHEMISTRY. vol. 29, no. 6, 13 February 1990, EASTON, PA US pages 1362 - 1367 GLOCKSHUBER R;MALIA M;PFITZINGER I;PLUCKTHUN A; 'A comparison of strategies to stabilize Immunoglobulin Fv-fragments' see page 1362, column 1, line 1 - page 1362, column 1, line 17 see page 1364, column 1, line 18 - page 1364, column 2, line 38 Y NATURE. vol. 349, 24 January 1991, LONDON GB pages 293 - 299 WINTER, G.; MILSTEIN, C.; 'Man-made antibodies' see page 295, column 2, line 14 - page 295, column 2, line 27 see page 297, column 1, line 34 - page 297, column 1, line 55; figure 2 Y NATURE. vol. 341, 12 October 1989, LONDON GB pages 544 - 546 WARD ES; GUSSOW D; GRIFFITHS AD; JONES PT; WINTER G; 'Binding activities of a repertoire of single immunoglobulin variable domains secreted from Escherichia coli'	20, 42,46 .10, 17-20
NATURE. Yol. 29, no. 6, 13 February 1990, EASTON, PA US pages 1362 - 1367 GLOCKSHUBER R;MALIA M;PFITZINGER I;PLUCKTHUN A; 'A comparison of strategies to stabilize Immunoglobulin Fv-fragments' see page 1362, column 1, line 1 - page 1362, column 1, line 17 see page 1364, column 1, line 18 - page 1364, column 2, line 38 Y NATURE. YOL. 349, 24 January 1991, LONDON GB pages 293 - 299 WINTER, G.; MILSTEIN, C.; 'Man-made antibodies' see page 295, column 2, line 14 - page 295, column 2, line 27 see page 297, column 1, line 34 - page 297, column 1, line 55; figure 2 Y NATURE. YOL. 341, 12 October 1989, LONDON GB pages 544 - 546 WARD ES;GUSSOW D;GRIFFITHS AD;JONES PT;WINTER G; 'Binding activities of a repertoire of single immunoglobulin variable domains secreted from Escherichia coli'	20, 42,46 .10, 17-20
vol. 349, 24 January 1991, LONDON GB pages 293 - 299 WINTER, G.; MILSTEIN, C.; 'Man-made antibodies' see page 295, column 2, line 14 - page 295, column 2, line 27 see page 297, column 1, line 34 - page 297, column 1, line 55; figure 2 NATURE. vol. 341, 12 October 1989, LONDON GB pages 544 - 546 WARD ES; GUSSOW D; GRIFFITHS AD; JONES PT; WINTER G; 'Binding activities of a repertoire of single immunoglobulin variable domains secreted from Escherichia coli'	17 -20
vol. 341, 12 October 1989, LONDON GB pages 544 - 546 WARD ES;GUSSOW D;GRIFFITHS AD;JONES PT;WINTER G; 'Binding activities of a repertoire of single immunoglobulin variable domains secreted from Escherichia coli'	12
see the whole document	
	,10, 17-20
-	
	·
	·

PC' JS 93/04338

Box 1 Observations where certain claims were found unscarchable (Continuation of item 1 of first sheet)
This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 32,34,36,38,40,50 are directed to a method of treatment of the human/animal body as well as diagnostic methods (Rule 39.1(iv) PCT) the search has been carried out and based on the alleged effects of the compound/composition. 2. Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a). Box 11 Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searches without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.;
Remark on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 9304338 SA 74117

This amer lists the patent family members relating to the patent documents cited in the above-mentioned international search report.

The members are as contained in the European Patent fixe EDP file on

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10/09/93

Patent document cited in search report	Publication date	Patent mem	family ber(s)	Publication date	
WO-A-8809344	01-12-88	AU-B- AU-A- AU-A- EP-A- JP-T- US-A- US-A-	612370 1804988 8579991 0318554 2500329 5132405 5091513	11-07-91 21-12-88 13-02-92 07-06-89 08-02-90 21-07-92 25-02-92	
	.* •				
	2				
			÷	•	
				:	
		·			
·	e e e e e e e e e e e e e e e e e e e		• • •		
		·			
		•			
			٠		
	•				

ORM POSTS

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82